

Channel Islands Cancer Report 2013

Incorporating data up to 2011

Produced by the Public Health England Knowledge and Intelligence Team
(South West), for the Guernsey and Jersey Medical Officers of Health

PUBLIC REPORT
(numbers less than 5 are suppressed for confidentiality)

Prepared by: Luke Hounsome, Principal Cancer Intelligence Analyst, Public Health England Knowledge and Intelligence Team (South West).
For queries relating to this document, please contact: +44 (0) 11 7970 6474.
Published .. 2013

The National Cancer Intelligence Network (NCIN) is a UK-wide partnership operated by Public Health England. The NCIN coordinates and develops analysis and intelligence to drive improvements in prevention, standards of cancer care and clinical outcomes for cancer patients.

Contents

Introduction	5
Methods.....	5
Data sources.....	5
Data quality.....	6
Methods.....	6
Populations	7
Key results.....	8
Guernsey	9
Incidence.....	9
Mortality	9
Jersey	10
Incidence.....	10
Mortality	11
All Cancers (C00-C97 excluding C44) - INCIDENCE.....	17
All Cancers (C00-C97 excluding C44) - MORTALITY	18
Bladder Cancer (C67) - INCIDENCE.....	19
Bladder Cancer (C67) - MORTALITY	20
Brain and CNS Cancer (C70-72) - INCIDENCE	21
Brain and CNS Cancer (C70-72) - MORTALITY.....	22
Breast Cancer (females) (C50) - INCIDENCE.....	23
Breast Cancer (females) (C50) - MORTALITY	24
Colorectal Cancer (C18-C21) - INCIDENCE	25
Colorectal Cancer (C18-C21) - MORTALITY	26
Head and Neck Cancer (C00-C14, C30-C32, C73) - INCIDENCE.....	27
Head and Neck Cancer (C00-C14, C30-C32, C73) - MORTALITY.....	28
Hepatobiliary Cancer (C22-C23) - INCIDENCE.....	29
Hepatobiliary Cancer (C22-C23) - MORTALITY.....	30
Kidney and Ureter Cancer (C64-C68) - INCIDENCE	31
Kidney and Ureter Cancer (C64-C68) - MORTALITY	32
Leukaemia (C91-C95) - INCIDENCE	33
Leukaemia (C91-C95) - MORTALITY	34
Lung Cancer (C33-C34) - INCIDENCE	35
Lung Cancer (C33-C34) - MORTALITY.....	36
Lymphoma (C81-C85, C96) - INCIDENCE.....	37
Lymphoma (C81-C85, C96) - MORTALITY	38
Male Urogenital Cancer excl. Prostate (C60-C63 excl. C61) - INCIDENCE	39
Male Urogenital Cancer excl. Prostate (C60-C63 excl. C61) - MORTALITY	40

Malignant Melanoma (C43) - INCIDENCE	41
Malignant Melanoma (C43) - MORTALITY	42
Other Cancers (other 'C' codes excl. C44) - INCIDENCE.....	43
Other Cancers (other 'C' codes excl. C44) - MORTALITY	44
Other Gynaecological Cancers (C51-C53, C57-C58) - INCIDENCE	45
Other Gynaecological Cancers (C51-C53, C57-C58) - MORTALITY.....	46
Other Skin Cancers (C44) - INCIDENCE.....	47
Other Skin Cancers (C44) - MORTALITY	48
Ovarian Cancer (C56) - INCIDENCE	49
Ovarian Cancer (C56) - MORTALITY	50
Prostate Cancer (C61) - INCIDENCE	51
Prostate Cancer (C61) - MORTALITY	52
Upper Gastrointestinal Cancers (C15-C16, C25) - INCIDENCE	53
Upper Gastrointestinal Cancers (C15-C16, C25) - MORTALITY	54
Uterine Cancer (C54-C55) - INCIDENCE	55
Uterine Cancer (C54-C55) - MORTALITY	56
Staging completeness	57
Appendix – ICD-10 codes	58

Introduction

Public Health England's Knowledge and Intelligence Team (KIT) South West (formerly the South West Public Health Observatory) analyses data on cancers diagnosed in residents of the Channel Islands, on behalf of their Public Health departments. As part of this process it produces an annual report on the basic statistics on cancer in the Channel Islands.

KIT South West hold cancer incidence data for the Channel Islands and the South West of England to 2011, plus data for England to 2010. Data on deaths from cancer and other causes are available to 2011 for all areas.

Methods

Data sources

Data on incidence are supplied by the health departments of Jersey and Guernsey from hospital records and pathology laboratories. Data from Guernsey are supplemented by data from the hospital Multi-Disciplinary Team (MDT), for staging of cancers. Guernsey figures include cancer deaths in residents of Alderney and incident cancers in residents of Alderney and Sark.

Cancers are coded according to the International Classification of Diseases v10 (ICD-10) coding system. The cancers are grouped according to the table below. A full list of ICD codes and descriptions is in the appendix. The data available are for 1999 to 2011 inclusive, except for data for the whole of England which is only available until 2010.

Cancer groups and ICD-10 codes

Cancer Site	ICD-10 codes
All Cancers excluding non-melanoma skin	C00-C97 excl. C44
Bladder	C67
Brain and CNS	C70-C72
Breast	C50
Colorectal	C18-C21
Head & neck	C00-C14, C30-C32, C73
Hepatobiliary	C22-C24
Kidney and ureter	C64-C66
Leukaemia	C91-C95
Lung	C33-C34
Lymphoma	C81-C85, C96
Male urogenital (excl. prostate)	C60-C63 excl. C61
Malignant melanoma	C43
Non-melanoma skin cancer (NMSC)	C44
Other cancer	Other Cs
Other gynaecological	C51-C53, C57-C58
Ovary	C56
Paediatric (0-19 years)	C00-C97 excl. C44
Prostate	C61
Upper gastrointestinal	C15-C16, C25
Uterus	C54-C55

Data on deaths are supplied from deaths registers on each island. The registered causes of death are processed by the Office for National Statistics (ONS) according to their rules to find a single underlying cause of death. This underlying cause is used to identify those dying from cancer. The data available are for 2001 to 2011 inclusive.

Data quality

The quality of cancer registration data in the UK is amongst the highest in the world, and a substantial amount of time is spent ensuring data are accurate. The introduction of electronic pathology feeds to the cancer registry has brought a higher level of accuracy in diagnosis to the registration process. However, the unique situation regarding registration of data for the Channel Islands does cause some difficulties. Because the health systems in Jersey and Guernsey do not use NHS number it is not possible to link patients who have been diagnosed or had previous cancers when living on the UK mainland, and subsequently moved to the Channel Islands, and vice versa. This means that some cancers which are technically recurrences of a previous cancer; cancers diagnosed many years previously and detected at death; and secondary cancers from previously diagnosed primary cancers, may be registered as if they were new cases. The recurrence rate of many cancers is low which makes this an uncommon occurrence, whilst better pathology reporting makes it easier to determine if cancers are secondary to another cancer, therefore this is likely to be a relatively small problem.

Data on deaths are based on those living in the Channel Islands at time of death and hence have none of the problems described above. The quality of data will only be affected by the accuracy of the death certificate completed by the doctor certifying the death.

Methods

The risk of developing or dying from cancer is strongly influenced by age and sex, and so the makeup of the population in an area will affect the number of cases/deaths. To allow a fair comparison between different areas, or the same area through time, directly age-standardised rates (ASRs) are calculated. This method looks at the occurrence of cancer/death in the population of interest, and calculates what the rate would be in a pre-determined population (known as the standard population), controlling for age. If the same standard population is used for all populations analysed, then the rates are directly comparable with each other to measure whether there are any differences of significance. Rates in this report are standardised to the 1976 European Standard Population.

When ASRs are reported, the 95% confidence intervals are also reported (Lower Confidence Interval: LCI, Upper Confidence Interval: UCI). The confidence interval is a statistical concept which defines the degree of variation in a rate which can be considered normal. Occurrence of, or death from, a disease depends on complicated relationships between a multitude of risk factors, so there are fluctuations over time or geography which do not necessarily reflect a real increase or decrease in a rate. As the population of interest becomes smaller, an increase or decrease of just one or two cases/deaths has a larger effect on the rate. Hence the smaller the population and the smaller the number of cases/deaths, the larger the confidence interval.

Generally speaking, if the confidence interval of two ASRs overlap it is not possible to say that there is a statistically significant difference between the area investigated and the comparison area. 95% confidence intervals mean that out of 100 comparisons/tests we could expect five to show a significant difference when none exists, or vice versa.

For some data, more sophisticated statistical tests are used. These report the likelihood that two rates are not significantly different (for example, that the observed differences could have occurred by chance) as a p-value. It is convention to accept a p-value of 0.05 or less as evidence that the rates are significantly different.

The maximum and minimum values in England are calculated for the former Government Office Regions (of which South West is one). The former GORs are: North East, North West, Yorkshire and the Humber, East Midland, West Midlands, East of England, London, South East and the South West.

Unless otherwise specified, counts and rates are for those aged 20 years and older.

Populations

Populations used for England and the South West are mid-year estimates supplied by the Office for National Statistics (ONS), based on the 2001 Census and taking into account births, deaths and migration.

Populations used for Guernsey are supplied by the Guernsey Health Department. They are estimated by Guernsey's central policy unit based on social security data and data from the education department.

Populations used for Jersey are supplied by the Jersey Health Department. They are end-of-year estimates produced by the States of Jersey Statistics Unit, who use a population model based on baseline figures from the 2011 Jersey Census and adjusted for population changes by administrative data, such as Social Security and Education department data, and migration information from the Census.

Key results

This section highlights those cancers which have a high or low rate compared to the South West or England. All comparisons are based on aggregated data for 2007-2011 (except England data which are for 2006-10). Risk factors for those cancers are listed.

In all these comparisons, the number of cases/deaths should be taken into account as well as the comparison of rates. As numbers for some cancers are small, a large relative incidence/mortality rate may only reflect a few extra cases/deaths each year and hence are clinically less important.

Where a rate is described as higher or lower, it means that it is statistically significant i.e. the differences seen are unlikely to be just a chance variation. If differences are likely to be only by chance, they are described as 'similar'.

Unless otherwise specified, counts and rates are for those aged 20 years and older.

In tables, '-' indicates that there were no cases/deaths in that time-period for that cancer. <5 indicates that there were less than 5 cases/deaths in that time-period for that cancer.

Top 10 incident cancers by age-standardised rate, for Guernsey (2007-2011), Jersey (2007-2011) South West (2007-2011) and England (2006-2010)

Guernsey		Jersey		South West		England	
Cancer Site	ASR	Cancer Site	ASR	Cancer Site	ASR	Cancer Site	ASR
NMSC	370.2	NMSC	361.7	NMSC	248.6	Breast	176.2
Breast	153.0	Breast	180.8	Breast	185.5	NMSC	160.5
Prostate	134.3	Prostate	175.9	Prostate	160.7	Prostate	148.1
Colorectal	66.2	Colorectal	65.0	Colorectal	68.8	Lung	65.9
Lung	59.9	Lung	62.7	Lung	54.6	Colorectal	65.6
Malignant Melanoma	50.9	Malignant Melanoma	47.9	Upper Gastrointestinal	35.8	Upper Gastrointestinal	38.7
Upper Gastrointestinal	41.5	Head and Neck	36.2	Malignant Melanoma	30.4	Uterus	27.5
Head and Neck	31.4	Upper Gastrointestinal	34.4	Uterus	29.0	Lymphoma	24.1
Bladder	31.2	Lymphoma	26.2	Lymphoma	26.1	Ovary	23.4
Lymphoma	28.5	Uterus	20.7	Ovary	24.4	Head and Neck	22.4

Note:

All cancers and 'other' cancers excluded from list. NMSC is non-melanoma skin cancer.

Top 10 cancer causes of death by age-standardised rate, for Guernsey , Jersey, South West and England (2007-2011)

Guernsey		Jersey		South West		England	
Cancer Site	ASR	Cancer Site	ASR	Cancer Site	ASR	Cancer Site	ASR
Lung	49.5	Lung	55.5	Lung	42.8	Lung	52.9
Prostate	44.2	Prostate	36.0	Prostate	34.8	Breast	35.7
Upper Gastrointestinal	34.6	Breast	32.9	Breast	34.5	Prostate	33.9
Colorectal	23.4	Upper Gastrointestinal	28.4	Upper Gastrointestinal	29.8	Upper Gastrointestinal	31.3
Breast	21.8	Colorectal	19.9	Colorectal	23.0	Colorectal	23.4
Ovary	14.1	Head and Neck	13.1	Ovary	12.6	Ovary	12.7
Brain and CNS	10.1	Ovary	12.5	Lymphoma	7.5	Lymphoma	7.5
Kidney and Ureter	10.1	Kidney and Ureter	10.6	Brain and CNS	7.5	Bladder	6.8
Bladder	8.5	Lymphoma	9.5	Leukaemia	6.7	Brain and CNS	6.7
Head and Neck	8.2	Hepatobiliary	8.9	Bladder	6.5	Hepatobiliary	6.7

Note:

All cancers and 'other' cancers excluded from list.

Guernsey

Incidence

The rate of breast cancer in Guernsey is lower than in the South West, but similar to the England average. The age-standardised rate (ASR) is 153 per 100,000. The rate in the South West is 186 per 100,000. The rate in Guernsey is similar to the lowest ASR in the England regions, which is 166 per 100,000. Around 45 new cases of breast cancer are diagnosed each year. Excessive alcohol intake, hormone replacement therapy and genetic/family links, and are some of the main risks for breast cancer.

The rate of prostate cancer in Guernsey is lower than in the South West, but similar to the England average. The ASR is 134 per 100,000. The rate in the South West is 160 per 100,000. The rate is similar to the lowest ASR in the England regions, which is 124 per 100,000. Around 40 new cases of prostate cancer are diagnosed each year. No major environmental/lifestyle risks for prostate cancer are known; risks are increased in those having a family member with the disease, and in black men.

The rate of malignant melanoma in Guernsey is higher than in the South West or England. The ASR is 51 per 100,000. The rate in the South West is 30 per 100,000, and in England is 22 per 100,000. The South West has the highest incidence of malignant melanoma in England. Around 27 new cases of malignant melanoma are diagnosed each year, and it is one of the most common cancers in those aged under 40. The major risk factor is UV exposure through sunlight or sunbeds.

The rate of bladder cancer in Guernsey is higher than in the South West or England. The ASR is 31 per 100,000. The rate in the South West is 15 per 100,000, and in England is 16 per 100,000. It is higher than the highest ASR in the England regions, which is 20 per 100,000. Around 20 new cases of bladder cancer are diagnosed each year. The main risk factors are smoking and industrial exposures.

The rate of head and neck cancers in Guernsey is higher than in the South West or England. The ASR is 31 per 100,000. The rate in the South West is 22 per 100,000, and in England is 22 per 100,000. It is similar to the highest ASR in the England regions, which is 26 per 100,000. Around 17 new cases of head and neck cancer are diagnosed each year. The main risk factors are smoking and excessive alcohol use, especially in combination.

The rate of non-melanoma skin cancers in Guernsey is higher than the *recorded* rate in the South West or England. However, care must be taken as not all parts of England record non-melanoma skin cancer comprehensively. The data registration in the Channel Islands and the South West is comparable, so we can be confident that the ASR is higher than in the South West. The ASR in Guernsey is 379 per 100,000 and in the South West is 249 per 100,000. Around 242 new cases in Guernsey are diagnosed each year. The risk factors are the same as those for malignant melanoma.

Mortality

The death rate from all cancers (excluding non-melanoma skin) in Guernsey is lower than the England average. The ASR is 219 per 100,000. The rate in England is 237 per 100,000. It is similar to the lowest ASR in the England regions, which is 221 per 100,000. Around 156 people in Guernsey die from cancer each year.

The death rate from prostate cancer in Guernsey is higher than the England average. The ASR is 44 per 100,000. The rate in England is 34 per 100,000. It is similar to the highest ASR in the England regions, which is 36 per 100,000. Around 15 men die from prostate cancer each year.

The death rate from breast cancer in Guernsey is lower than in the South West and in England as a whole. The ASR is 22 per 100,000. The rate in the South West is 35 per 100,000 and in England is 36 per 100,000. The South West has the lowest mortality from breast cancer in England. Around 8 women in Guernsey die from breast cancer each year

Jersey

Incidence

The rate of all cancers in Jersey is similar to the South West, but higher than the England average. The age-standardised rate (ASR) is 571 per 100,000. The rate in England is 544 per 100,000. Non-melanoma skin cancers are excluded from this comparison due to data issues (see below). The ASR in Jersey is comparable to the highest ASR in the England regions; which is 578 per 100,000. Around 488 new cancers (excluding non-melanoma skin cancer) are diagnosed each year in Jersey.

The rate of prostate cancer in Jersey is similar to the South West, but higher than the England average. The ASR is 176 per 100,000. The rate in England is 148 per 100,000. The South West has the highest incidence of prostate cancer in England. Around 71 new cases of prostate cancer are diagnosed each year in Jersey. No major environmental/lifestyle risks for prostate cancer are known; risks are increased in those having a family member with the disease, or in black men.

The rate of lung cancer in Jersey is higher than in the South West, but similar to the England average. The ASR is 63 per 100,000. The rate in the South West is 55 per 100,000. It is lower than the highest ASR in the England regions, which is 96 per 100,000. Around 54 new cases of lung cancer are diagnosed each year. Smoking is responsible for about 85% of lung cancers.

The rate of malignant melanoma in Jersey is higher than in the South West or England. The ASR is 48 per 100,000. The rate in the South West is 30 per 100,000, and in England is 22 per 100,000. The South West has the highest incidence of malignant melanoma in England. Around 39 new cases of malignant melanoma are diagnosed each year, and it is one of the most common cancers in those aged under 40. The major risk factor is UV exposure through sunlight or sunbeds.

The rate of head and neck cancers in Jersey is higher than in the South West or England. The ASR is 36 per 100,000. The rate in the South West is 22 per 100,000, and in England is 22 per 100,000. It is higher than the highest ASR in the England regions, which is 26 per 100,000. Around 28 new cases of head and neck cancer are diagnosed each year. The main risk factors are smoking and excessive alcohol use, especially when in combination.

The rate of leukaemia in Jersey is similar to the South West, but higher than the England average. The ASR is 18 per 100,000. The rate in England is 13 per 100,000. The ASR in Jersey is similar to the highest ASR in the England regions; which is 16 per 100,000. Around 14 new cases of leukaemia are diagnosed each year in Jersey. The risk factors for leukaemia vary depending on the exact type but can include smoking, exposure to chemicals and exposure to radiation far in excess of background levels.

The rate of kidney and ureter cancer in Jersey is lower than in the South West, but similar to the England average. The ASR is 12 per 100,000. The rate in the South West is 17 per 100,000. It is similar to the lowest ASR in the England regions, which is 13 per 100,000. Around 10 new cases of kidney and ureter cancer are diagnosed each year in Jersey. The risk of kidney cancer is increased by being overweight or obese, by smoking and by certain genetic conditions.

The rate of uterine cancer in Jersey is lower than in the South West, but similar to the England average. The ASR is 21 per 100,000. The rate in the South West is 29 per 100,000. It is similar to the lowest ASR in the England regions, which is 20 per 100,000. Around 9 new cases of uterine cancer are diagnosed each year in Jersey. The risk of uterine cancer is increased by being overweight or obese, but decreased by having children and/or by use of the contraceptive pill.

The rate of ovarian cancer in Jersey is lower than in the South West, but similar to the England average. The ASR is 17 per 100,000. The rate in the South West is 24 per 100,000. It is lower than the lowest ASR in the England regions, which is 24 per 100,000. Around 8 new cases of ovarian cancer are diagnosed each year in Jersey. No major environmental/lifestyle risks for ovarian cancer have been well demonstrated, but there are genetic risks.

The rate of non-melanoma skin cancers in Jersey is higher than the *recorded* rate in the South West or England. However, care must be taken as not all parts of England record non-melanoma comprehensively. The data registration in the Channel Islands and the South West is comparable, so we can be confident that the ASR is higher than in the South West. The ASR in Jersey is 362 per 100,000 and the rate in the South West is 249 per 100,000. Around 318 new cases are diagnosed each year. The risk factors are the same as those for malignant melanoma

Mortality

The death rate for lung cancer in Jersey is higher than in the South West but similar to the England average. The age-standardised rate (ASR) is 56 per 100,000. The rate in the South West is 43 per 100,000. It is lower than the highest ASR in the England regions, which is 77 per 100,000. Around 48 people in Jersey die from lung cancer each year. The high death rate is likely to be reflective of the high incidence.

The death rate for head and neck cancers in Jersey is higher than in the South West and England. The ASR is 13 per 100,000. The rate in the South West is 5 per 100,000, and in England is 6 per 100,000. It is higher than the highest ASR in the England regions, which is 7 per 100,000. Around 8 people in Jersey die from a head and neck cancer each year. The high death rate is likely to be reflective of the high incidence.

The death rate for hepatobiliary cancer in Jersey is higher than in the South West but similar to the England average. The ASR is 9 per 100,000. The rate in the South West is 6 per 100,000. It is similar to the highest ASR in the England regions, which is 8 per 100,000. Around 8 people in Jersey die from hepatobiliary cancer each year.

The death rate for kidney and ureter cancers in Jersey is higher than in the South West and England. The ASR is 11 per 100,000. The rate in the South West is 6 per 100,000, and in England is 6 per 100,000. It is higher than the highest ASR in the England regions, which is 7 per 100,000. Around 7 people in Jersey die from a kidney or ureter cancer each year. The high death rate is in contrast to the low incidence.

The death rate for malignant melanoma in Jersey is similar to the South West but higher than the England average. The ASR is 8 per 100,000. The rate in England is 4 per 100,000. It is similar to the highest ASR in the England regions, which is 5 per 100,000. Around 4 people in Jersey die from malignant melanoma each year. The high death rate is likely to be reflective of the high incidence.

The death rate from 'other' cancers in Jersey is lower than the South West and the England average. The ASR is 24 per 100,000. The rate in the South West is 31 per 100,000, and in England is 34 per 100,000. It is lower than the lowest ASR in the England regions, which is 32 per 100,000. Around 21 people in Jersey die from 'other' cancers each year.

The death rate from 'other gynaecological' cancers in Jersey is lower than the South West and the England average. The ASR is 2 per 100,000. The rate in the South West is 5 per 100,000, and in England is 5 per 100,000. It is similar to the lowest ASR in the England regions, which is 4 per 100,000. Only 1 women in Jersey dies from 'other gynaecological' cancers each year, on average, so these figures should be interpreted in that context.

Age-standardised incidence rates (per 100,000 population) for Guernsey (2007-2011), South West (2007-2011), and England (2006-2010)

Age Group	Cancer Site	Expected cases*	Actual cases	Guernsey					South West					England					Guernsey compared to England		Guernsey compared to Jersey			
				ASR	ASR LCI	ASR UCI	Cases	ASR	ASR LCI	ASR UCI	p-value	difference	Cases	ASR	ASR LCI	ASR UCI	p-value	difference	Cases	ASR	ASR LCI	ASR UCI	p-value	difference
20+	All Cancers excl. C44	1,707	1,740	568.7	531.5	585.8	154,959	557.3	554.4	560.3	0.923	-	1,300,430	543.6	542.6	544.5	0.277	-	0.249	-				
20+	Bladder	57	99	31.2	24.3	38.1	5,024	15.4	14.9	15.9	0.000	higher	44,079	16.3	16.2	16.5	0.000	higher	0.000	higher				
20+	Brain and CNS	25	21	9.8	5.3	14.4	2,299	9.5	9.1	9.9	0.883	-	19,049	8.8	8.6	8.9	0.640	-	0.172	-				
20+	Breast	265	225	153.0	132.3	173.7	23,922	185.5	183.0	188.0	0.002	lower	200,603	176.2	175.4	177.0	0.029	-	0.023	lower				
20+	Colorectal	218	209	66.2	56.8	75.6	20,594	68.8	67.8	69.8	0.585	-	165,685	65.6	65.3	66.0	0.907	-	0.423	-				
20+	Head and Neck	64	85	31.4	24.4	38.4	5,179	21.5	20.8	22.1	0.006	higher	47,955	22.4	22.2	22.6	0.011	higher	0.153	-				
20+	Hepatobiliary	30	19	9.3	4.7	13.9	2,642	8.9	8.6	9.3	0.881	-	22,611	9.0	8.9	9.1	0.895	-	0.420	-				
20+	Kidney and Ureter	48	38	15.7	10.1	21.3	4,685	16.9	16.4	17.4	0.685	-	36,108	15.2	15.1	15.4	0.866	-	0.134	-				
20+	Leukaemia	36	50	18.2	12.4	23.9	4,032	13.8	13.4	14.3	0.138	-	32,132	13.0	12.9	13.2	0.078	-	0.442	-				
20+	Lung	218	189	59.9	51.1	68.8	16,515	54.6	53.7	55.5	0.243	-	167,244	65.9	65.6	66.2	0.187	-	0.314	-				
20+	Lymphoma	73	82	28.5	21.8	35.1	6,909	26.1	25.5	26.8	0.492	-	55,378	24.1	23.9	24.4	0.202	-	0.290	-				
20+	Male Urogenital (excl. Prostate)	14	13	11.0	5.0	17.0	1,311	14.0	13.3	14.8	0.322	-	11,047	11.8	11.5	12.0	0.802	-	0.399	-				
20+	Malignant Melanoma	65	136	50.9	41.4	60.4	7,424	30.4	29.7	31.2	0.000	higher	48,457	22.4	22.2	22.6	0.000	higher	0.290	-				
20+	NMSC	547	1,209	370.2	348.5	392.0	73,293	248.6	246.7	250.5	0.000	higher	407,020	160.5	160.0	161.0	0.000	higher	0.272	-				
20+	Other Cancers	133	143	43.2	35.7	50.7	11,791	39.4	38.6	40.1	0.313	-	104,329	40.9	40.7	41.2	0.544	-	0.043	higher				
20+	Other Gynaecological	25	30	21.3	13.3	29.3	2,301	20.1	19.2	21.0	0.766	-	19,289	17.6	17.3	17.8	0.356	-	0.169	-				
20+	Ovary	36	30	18.2	11.3	25.1	3,335	24.4	23.5	25.3	0.079	-	27,720	23.4	23.1	23.6	0.141	-	0.426	-				
20+	Prostate	219	200	134.3	115.5	153.2	21,916	160.7	158.5	162.9	0.007	lower	166,840	148.1	147.3	148.8	0.155	-	0.001	lower				
20+	Upper Gastrointestinal	130	131	41.5	33.9	49.0	11,066	35.8	35.1	36.5	0.143	-	99,611	38.7	38.5	39.0	0.478	-	0.058	-				
20+	Uterus	43	40	25.6	17.3	34.0	4,014	29.0	28.1	30.0	0.427	-	32,293	27.5	27.2	27.8	0.663	-	0.161	-				

Note:

ASR – Directly Age Standardised Incidence Rate (significance testing is at 95% significance level); LCI – 95% Lower Confidence Interval; UCI – 95% Upper Confidence Interval; Higher – Rate is statistically significantly higher than the comparator; Lower – Rate is statistically significantly lower than the comparator.

NMSC is non-melanoma skin cancer.

*Expected cases based on the England average.

Age-standardised incidence rates (per 100,000 population) for Jersey (2007-2011), South West (2007-2011), and England (2006-2010)

Age Group	Cancer Site	Jersey						South West						England						Jersey compared to South West		Jersey compared to England		Jersey compared to Guernsey						
		Expected cases*	Actual cases	ASR	ASR LCI	ASR UCI	Cases	ASR	ASR LCI	ASR UCI	Cases	ASR	ASR LCI	ASR UCI	Cases	ASR	ASR LCI	ASR UCI	Cases	ASR	ASR LCI	ASR UCI	p-value	difference	p-value	difference	p-value	difference	p-value	difference
20+	All Cancers excl. C44	2,359	2,438	570.7	547.5	593.8	154,959	557.3	544.4	560.3	1,300,430	543.6	542.6	544.5	1,300,430	543.6	542.6	544.5	1,300,430	543.6	542.6	544.5	0.262	0.022	0.249	-	0.022	higher	0.249	-
20+	Bladder	77	78	17.9	13.4	22.4	5,024	15.4	14.9	15.9	44,079	16.3	16.2	16.5	44,079	16.3	16.2	16.5	44,079	16.3	16.2	16.5	0.499	0.499	0.000	lower	0.499	-	0.000	lower
20+	Brain and CNS	36	27	7.5	4.5	10.4	2,299	9.5	9.1	9.9	19,049	8.8	8.6	8.9	19,049	8.8	8.6	8.9	19,049	8.8	8.6	8.9	0.185	0.396	0.172	-	0.396	-	0.172	-
20+	Breast	376	387	180.8	162.3	199.3	23,922	185.5	183.0	188.0	200,603	176.2	175.4	177.0	200,603	176.2	175.4	177.0	200,603	176.2	175.4	177.0	0.623	0.622	0.023	higher	0.622	-	0.023	higher
20+	Colorectal	296	281	65.0	57.1	73.0	20,594	68.8	67.8	69.8	68,585	65.6	65.3	66.0	68,585	65.6	65.3	66.0	68,585	65.6	65.3	66.0	0.351	0.880	0.423	-	0.880	-	0.423	-
20+	Head and Neck	92	141	36.2	30.1	42.2	5,179	21.5	20.8	22.1	47,955	22.4	22.2	22.6	47,955	22.4	22.2	22.6	47,955	22.4	22.2	22.6	0.000	0.000	0.420	-	0.000	higher	0.420	-
20+	Hepatobiliary	41	40	9.8	6.6	13.1	2,642	8.9	8.6	9.3	22,611	9.0	8.9	9.1	22,611	9.0	8.9	9.1	22,611	9.0	8.9	9.1	0.596	0.611	0.153	-	0.611	-	0.153	-
20+	Kidney and Ureter	66	48	12.4	8.6	16.1	4,685	16.9	16.4	17.4	36,108	15.2	15.1	15.4	36,108	15.2	15.1	15.4	36,108	15.2	15.1	15.4	0.021	0.137	0.134	-	0.137	-	0.134	-
20+	Leukaemia	50	72	17.7	13.3	22.1	4,032	13.8	13.4	14.3	32,132	13.0	12.9	13.2	32,132	13.0	12.9	13.2	32,132	13.0	12.9	13.2	0.086	0.037	0.442	-	0.037	higher	0.442	-
20+	Lung	295	270	62.7	55.0	70.4	16,515	54.6	53.7	55.5	167,244	65.9	65.6	66.2	167,244	65.9	65.6	66.2	167,244	65.9	65.6	66.2	0.040	0.423	0.314	-	0.423	-	0.314	-
20+	Lymphoma	103	100	26.2	20.8	31.6	6,909	26.1	25.5	26.8	55,378	24.1	23.9	24.4	55,378	24.1	23.9	24.4	55,378	24.1	23.9	24.4	0.977	0.456	0.290	-	0.456	-	0.290	-
20+	Male Urogenital (excl. Prostate)	22	22	12.0	7.0	17.1	1,311	14.0	13.3	14.8	11,047	11.8	11.5	12.0	11,047	11.8	11.5	12.0	11,047	11.8	11.5	12.0	0.435	0.920	0.399	-	0.920	-	0.399	-
20+	Malignant Melanoma	92	194	47.9	41.0	54.8	7,424	30.4	29.7	31.2	48,457	22.4	22.2	22.6	48,457	22.4	22.2	22.6	48,457	22.4	22.2	22.6	0.000	0.000	0.290	-	0.000	higher	0.290	-
20+	NMISC	747	1,590	361.7	343.5	380.0	73,293	248.6	246.7	250.5	407,020	160.5	160.0	161.0	407,020	160.5	160.0	161.0	407,020	160.5	160.0	161.0	0.000	0.000	0.272	-	0.000	higher	0.272	-
20+	Other Cancers	182	149	35.4	29.5	41.3	11,791	39.4	38.6	40.1	104,329	40.9	40.7	41.2	104,329	40.9	40.7	41.2	104,329	40.9	40.7	41.2	0.190	0.066	0.043	lower	0.066	-	0.043	lower
20+	Other Gynaecological	37	34	16.8	11.0	22.6	2,301	20.1	19.2	21.0	19,289	17.6	17.3	17.8	19,289	17.6	17.3	17.8	19,289	17.6	17.3	17.8	0.265	0.789	0.169	-	0.789	-	0.169	-
20+	Ovary	50	40	17.4	11.8	23.0	3,335	24.4	23.5	25.3	27,720	23.4	23.1	23.6	27,720	23.4	23.1	23.6	27,720	23.4	23.1	23.6	0.015	0.036	0.426	-	0.036	-	0.426	-
20+	Prostate	299	355	175.9	157.4	194.5	21,916	160.7	158.5	162.9	166,840	148.1	147.3	148.8	166,840	148.1	147.3	148.8	166,840	148.1	147.3	148.8	0.109	0.003	0.001	higher	0.003	higher	0.001	higher
20+	Upper Gastrointestinal	176	153	34.4	28.8	40.0	11,066	35.8	35.1	36.5	99,611	38.7	38.5	39.0	99,611	38.7	38.5	39.0	99,611	38.7	38.5	39.0	0.625	0.130	0.058	-	0.130	-	0.058	-
20+	Uterus	59	47	20.7	14.6	26.9	4,014	29.0	28.1	30.0	32,293	27.5	27.2	27.8	32,293	27.5	27.2	27.8	32,293	27.5	27.2	27.8	0.009	0.031	0.161	-	0.031	-	0.161	-

Note:

ASR – Directly Age Standardised Incidence Rate (significance testing is at 95% significance level); LCI – 95% Lower Confidence Interval; UCI – 95% Upper Confidence Interval; Higher – Rate is statistically significantly higher than the comparator; Lower – Rate is statistically significantly lower than the comparator.

NMISC is non-melanoma skin cancer.

*Expected cases based on the England average.

Age-standardised mortality rates (per 100,000 population) for Guernsey, South West and England, 2007-2011

Age Group	Cancer Site	Guernsey					South West					England					Guernsey compared to England		Guernsey compared to Jersey	
		Expected deaths*	Actual deaths	ASR	ASR LCI	ASR UCI	Deaths	ASR	ASR LCI	ASR UCI	p-value	difference	Deaths	ASR	ASR LCI	ASR UCI	p-value	difference	p-value	difference
20+	All Cancers excl. C44	830	780	218.8	202.7	235.0	71,466	221.4	219.7	223.1	0.758	-	639,409	237.0	236.3	237.6	0.028	lower	0.326	-
20+	Bladder	27	29	8.5	4.7	12.2	2,421	6.5	6.2	6.7	0.293	-	20,816	6.8	6.7	6.9	0.384	-	0.101	-
20+	Brain and CNS	20	20	10.1	5.3	15.0	1,895	7.5	7.1	7.8	0.274	-	15,216	6.7	6.6	6.8	0.164	-	0.280	-
20+	Breast	65	39	21.8	14.4	29.2	5,540	34.5	33.5	35.6	0.001	lower	49,032	35.7	35.4	36.1	0.000	lower	0.017	lower
20+	Colorectal	85	67	23.4	16.8	30.0	7,700	23.0	22.5	23.6	0.144	-	65,329	23.4	23.2	23.6	0.993	-	0.158	-
20+	Head and Neck	18	17	8.2	4.0	12.4	1,418	5.0	4.7	5.3	0.144	-	13,817	5.7	5.6	5.8	0.250	-	0.054	-
20+	Hepatobiliary	22	11	6.1	2.3	9.8	1,837	5.9	5.6	6.2	0.917	-	17,627	6.7	6.6	6.8	0.751	-	0.132	-
20+	Kidney and Ureter	20	23	10.1	5.5	14.7	1,827	5.8	5.5	6.1	0.070	-	15,584	6.0	5.9	6.1	0.079	-	0.422	-
20+	Leukaemia	24	21	7.3	3.7	10.9	2,218	6.7	6.4	7.0	0.733	-	18,283	6.6	6.5	6.7	0.708	-	0.481	-
20+	Lung	182	168	49.5	41.7	57.3	13,387	42.8	42.0	43.6	0.095	-	139,630	52.9	52.6	53.2	0.392	-	0.129	-
20+	Lymphoma	26	15	6.8	2.7	10.9	2,410	7.5	7.1	7.8	0.748	-	20,078	7.5	7.4	7.6	0.751	-	0.142	-
20+	Male Urogenital (excl. Prostate)	1	-	-	-	-	86	0.7	0.5	0.9	-	-	740	0.7	0.6	0.7	-	-	-	-
20+	Malignant Melanoma	11	13	5.5	2.0	9.0	1,282	4.5	4.2	4.7	0.557	-	8,832	3.6	3.6	3.7	0.290	-	0.152	-
20+	NMSC	3	9	2.2	0.6	3.9	293	0.8	0.7	0.9	0.073	-	2,174	0.7	0.6	0.7	0.057	-	0.407	-
20+	Other Cancers	109	129	34.6	28.2	41.1	10,548	31.2	30.6	31.8	0.302	-	95,158	34.1	33.9	34.4	0.887	-	0.002	higher
20+	Other Gynaecological	8	5	2.6	0.1	5.1	711	4.8	4.4	5.2	0.093	-	6,238	4.7	4.6	4.8	0.108	-	0.357	-
20+	Ovary	22	26	14.1	8.2	20.0	2,046	12.6	12.0	13.2	0.615	-	17,229	12.7	12.5	12.9	0.644	-	0.325	-
20+	Prostate	56	74	44.2	34.0	54.5	5,627	34.8	33.8	35.7	0.070	-	44,208	33.9	33.5	34.2	0.046	higher	0.101	-
20+	Upper Gastrointestinal	109	113	34.6	27.4	41.7	9,608	29.8	29.1	30.4	0.190	-	84,133	31.3	31.1	31.5	0.369	-	0.064	-
20+	Uterus	9	10	5.5	1.8	9.3	925	5.3	5.0	5.7	0.912	-	7,459	5.2	5.0	5.3	0.837	-	0.230	-

Note:

ASR – Directly Age Standardised Mortality Rate (significance testing is at 95% significance level); LCI – 95% Lower Confidence Interval; UCI – 95% Upper Confidence Interval;

Higher – Rate is statistically significantly higher than the comparator; Lower – Rate is statistically significantly lower than the comparator.

NMSC is non-melanoma skin cancer.

*Expected cases based on the England average.

Age-standardised mortality rates (per 100,000 population) for Jersey, South West and England, 2007-2011

Age Group	Cancer Site	Jersey				South West				Jersey compared to South West				England				Jersey compared to England		Jersey compared to Guernsey	
		Expected deaths	Actual deaths	ASR	ASR LCI	ASR UCI	Deaths	ASR	ASR LCI	ASR UCI	p-value	difference	Deaths	ASR	ASR LCI	ASR UCI	p-value	difference	p-value	difference	
																					ASR
20+	All Cancers excl. C44	1118	1,020	223.6	209.4	237.7	71,466	221.4	219.7	223.1	0.764	639,409	237.0	236.3	237.6	0.064	-	-	0.326		
20+	Bladder	35	27	6.0	3.4	8.6	2,421	6.5	6.2	6.7	0.748	20,816	6.8	6.7	6.9	0.559	-	-	0.101		
20+	Brain and CNS	28	24	8.5	4.8	12.2	1,895	7.5	7.1	7.8	0.579	15,216	6.7	6.6	6.8	0.348	-	-	0.280		
20+	Breast	88	82	32.9	25.4	40.4	5,540	34.5	33.5	35.6	0.676	49,032	35.7	35.4	36.1	0.464	-	-	0.017		
20+	Colorectal	114	86	19.9	15.3	24.4	7,700	23.0	22.5	23.6	0.178	65,329	23.4	23.2	23.6	0.126	-	-	0.158		
20+	Head and Neck	25	39	13.1	8.8	17.3	1,418	5.0	4.7	5.3	0.000	13,817	5.7	5.6	5.8	0.001	higher	-	0.054		
20+	Hepatobiliary	30	39	8.9	6.0	11.7	1,837	5.9	5.6	6.2	0.040	17,627	6.7	6.6	6.8	0.132	-	-	0.132		
20+	Kidney and Ureter	27	33	10.6	6.6	14.7	1,827	5.8	5.5	6.1	0.021	15,584	6.0	5.9	6.1	0.025	higher	-	0.422		
20+	Leukaemia	32	29	7.4	4.3	10.5	2,218	6.7	6.4	7.0	0.645	18,283	6.6	6.5	6.7	0.617	-	-	0.481		
20+	Lung	245	242	55.5	48.3	62.7	13,387	42.8	42.0	43.6	0.001	139,630	52.9	52.6	53.2	0.491	-	-	0.129		
20+	Lymphoma	35	33	9.5	5.9	13.1	2,410	7.5	7.1	7.8	0.277	20,078	7.5	7.4	7.6	0.273	-	-	0.142		
20+	Male Urterial (excl. Prostate)	1	<5	1.1	-0.4	2.5	86	0.7	0.5	0.9	0.640	740	0.7	0.6	0.7	0.628	-	-	-		
20+	Malignant Melanoma	16	22	7.9	4.3	11.4	1,262	4.5	4.2	4.7	0.060	8,832	3.6	3.6	3.7	0.019	higher	-	0.152		
20+	NMSC	4	8	2.5	0.6	4.4	293	0.8	0.7	0.9	0.074	2,174	0.7	0.6	0.7	0.060	-	-	0.407		
20+	Other Cancers	146	107	24.0	19.2	28.7	10,548	31.2	30.6	31.8	0.003	95,158	34.1	33.9	34.4	0.000	lower	-	0.002		
20+	Other Gynaecological	11	6	2.1	0.3	3.9	711	4.8	4.4	5.2	0.004	6,238	4.7	4.6	4.8	0.004	lower	-	0.357		
20+	Ovary	30	32	12.5	8.0	17.1	2,046	12.6	12.0	13.2	0.976	17,229	12.7	12.5	12.9	0.932	-	-	0.325		
20+	Prostate	75	78	36.0	27.9	44.1	5,627	34.8	33.8	35.7	0.771	44,208	33.9	33.5	34.2	0.810	-	-	0.101		
20+	Upper Gastrointestinal	147	127	28.4	23.2	33.5	9,608	29.8	29.1	30.4	0.599	84,133	31.3	31.1	31.5	0.268	-	-	0.064		
20+	Uterus	13	12	4.0	1.7	6.4	925	5.3	5.0	5.7	0.295	7,459	5.2	5.0	5.3	0.362	-	-	0.230		

Note:

ASR – Directly Age Standardised Mortality Rate (significance testing is at 95% significance level); LCI – 95% Lower Confidence Interval; UCI – 95% Upper Confidence Interval;


Higher – Rate is statistically significantly higher than the comparator; Lower – Rate is statistically significantly lower than the comparator.

NMSC is non-melanoma skin cancer.


*Expected cases based on the England average.

All Cancers (C00-C97 excluding C44) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	549	530.1	485.5	574.8	574	471.0	430.9	511.1	415	569.8	514.5	625.2	398	461.6	413.5	509.7
2000-2002	541	521.5	477.2	565.7	581	476.2	436.0	516.5	433	597.8	540.9	654.7	414	475.2	426.5	523.8
2001-2003	607	582.4	535.8	629.1	600	494.7	453.6	535.8	453	632.7	573.8	691.5	429	502.3	451.9	552.7
2002-2004	702	667.5	617.8	717.3	646	526.5	484.3	568.7	483	670.6	610.2	731.1	474	529.4	478.5	580.3
2003-2005	741	694.2	643.8	744.7	659	531.3	489.0	573.6	488	676.6	615.9	737.3	507	576.7	523.4	630.1
2004-2006	803	739.1	687.4	790.7	663	521.6	480.1	563.0	457	603.5	547.7	659.4	489	534.7	484.5	585.0
2005-2007	800	710.2	660.5	759.9	680	523.2	482.3	564.2	476	601.8	547.1	656.4	457	497.1	448.9	545.3
2006-2008	840	720.4	671.2	769.6	677	509.4	469.5	549.2	494	586.2	533.8	638.6	452	472.4	426.1	518.6
2007-2009	775	644.0	598.1	689.9	677	504.0	464.6	543.4	589	683.1	627.2	739.0	482	492.5	445.8	539.2
2008-2010	779	641.8	596.2	687.4	671	501.4	462.2	540.6	580	657.7	603.4	712.1	488	501.9	454.8	549.0
2009-2011	779	643.4	597.7	689.1	661	497.1	458.0	536.1	556	625.4	572.7	678.2	511	525.9	477.8	573.9

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	68.8	69.8	67.0	68.8
Median	70	71	68	70
5th percentile	43	44	42	44
95th percentile	88	89	87	88

All Cancers (C00-C97 excluding C44) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	198	190.5	163.8	217.2	172	131.2	110.7	151.8	248	328.6	287.3	370.0	193	183.2	155.0	211.3
2002-2004	255	243.1	213.1	273.2	235	174.7	151.3	198.2	243	322.1	281.2	363.0	220	203.4	174.0	232.7
2003-2005	277	258.7	227.9	289.4	239	179.4	155.5	203.4	248	327.3	286.1	368.5	232	220.4	189.5	251.3
2004-2006	295	267.4	236.6	298.3	268	195.1	170.5	219.7	231	291.5	253.5	329.5	226	209.2	179.7	238.7
2005-2007	335	292.6	261.0	324.3	278	204.2	179.1	229.3	230	274.9	238.9	310.9	205	182.9	155.8	210.0
2006-2008	332	279.8	249.3	310.3	251	176.5	153.6	199.3	212	241.2	208.3	274.2	199	173.6	147.4	199.8
2007-2009	314	257.5	228.6	286.4	238	165.9	143.9	187.9	255	279.8	244.9	314.7	216	182.5	155.9	209.1
2008-2010	341	275.9	246.2	305.6	268	173.6	151.6	195.5	270	288.5	253.5	323.4	203	169.6	144.2	195.0
2009-2011	366	292.8	262.4	323.2	302	193.5	170.4	216.6	287	299.7	264.5	334.9	204	167.5	142.3	192.6

1-year survival from all cancers in males is rising in all areas. 5-year survival is rising in all areas.

1-year survival from all cancers in females is rising in all areas. 5-year survival is rising, except in Jersey where there is no evidence of change.


1-year survival in Guernsey, Jersey and the South West is higher than in England, for both males and females.

5-year survival for males in Guernsey is similar to England - in Jersey and the South West it is higher than in England.


5-year survival for females in Guernsey, Jersey and the South West is higher than in England.

Bladder Cancer (C67) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	40	39.0	26.8	51.1	13	9.5	4.1	14.9	40	53.9	37.1	70.8	13	10.8	4.5	17.1
2000-2002	34	33.3	22.0	44.5	16	11.9	5.8	18.0	35	48.1	32.0	64.1	7	7.6	1.5	13.6
2001-2003	34	32.1	21.2	43.0	17	13.0	6.6	19.5	36	51.2	34.4	68.0	5	4.8	0.2	9.3
2002-2004	39	36.0	24.6	47.4	18	12.4	6.3	18.5	34	49.0	32.4	65.5	9	9.6	2.9	16.4
2003-2005	34	30.4	20.1	40.7	15	9.2	4.1	14.2	35	49.1	32.7	65.4	11	10.9	4.1	17.8
2004-2006	33	29.6	19.4	39.7	15	8.0	3.6	12.4	33	41.6	27.2	56.0	18	15.8	8.0	23.6
2005-2007	28	23.5	14.6	32.3	14	7.4	3.3	11.5	40	46.7	32.0	61.4	20	15.0	7.9	22.1
2006-2008	42	33.6	23.3	43.9	14	8.0	3.5	12.5	35	38.9	25.7	52.0	23	16.2	9.2	23.3
2007-2009	40	29.9	20.5	39.3	11	6.2	2.2	10.1	37	42.1	28.3	55.8	26	21.7	12.8	30.5
2008-2010	42	30.8	21.4	40.2	7	4.1	0.9	7.2	34	38.1	25.0	51.1	24	20.1	11.6	28.6
2009-2011	32	23.5	15.3	31.7	10	5.5	1.9	9.2	38	40.9	27.6	54.1	20	16.5	8.9	24.2

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	74.1	74.9	75.7	73.7
Median	75	76	78	75
5th percentile	54	55	58	54
95th percentile	90	90	89	88

Bladder Cancer (C67) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	13	12.1	5.5	18.7	<5	1.1	-0.5	2.7	9	12.1	4.1	20.1	8	5.5	1.5	9.5
2002-2004	16	14.8	7.5	22.2	6	3.3	0.5	6.1	8	10.8	3.3	18.3	6	4.3	0.7	8.0
2003-2005	20	18.3	10.2	26.5	8	4.8	1.2	8.3	9	11.8	4.0	19.6	8	5.2	1.5	9.0
2004-2006	18	16.5	8.8	24.2	12	6.8	2.7	10.9	10	11.5	4.3	18.7	7	4.0	1.0	7.0
2005-2007	18	15.4	8.2	22.6	9	5.6	1.7	9.4	11	11.0	4.4	17.7	8	4.4	1.3	7.5
2006-2008	12	9.5	4.1	14.9	8	4.7	1.2	8.1	10	9.9	3.7	16.2	6	3.2	0.6	5.9
2007-2009	5	3.8	0.4	7.1	<5	2.6	-0.1	5.3	10	10.0	3.7	16.3	7	4.0	0.9	7.0
2008-2010	8	5.8	1.8	9.9	5	2.9	0.2	5.6	13	13.4	6.0	20.7	7	5.2	1.0	9.5
2009-2011	15	11.2	5.5	16.9	5	2.5	0.2	4.8	11	10.6	4.3	16.9	6	4.7	0.7	8.8

1-year survival from bladder cancer in males is rising in Jersey, and unchanged elsewhere. 5-year survival is rising in Jersey, and falling elsewhere.


1-year survival from bladder cancer in females is falling in Jersey, and unchanged elsewhere. 5-year survival is falling in Jersey and the South West, and unchanged elsewhere.

1-year and 5-year survival for bladder cancer in males is similar to England in Guernsey, Jersey and the South West.


1-year survival for bladder cancer in females is similar to England in Guernsey, Jersey and the South West. 5-year survival for bladder cancer in females is similar to England in Guernsey and Jersey but higher than England in the South West.

Brain and CNS Cancer (C70-72) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	9	9.0	3.1	14.9	5	4.3	0.4	8.3	6	8.9	1.8	16.1	<5	2.5	0	6.1
2000-2002	9	9.4	3.2	15.5	8	7.2	2.2	12.3	5	7.3	0.9	13.7	5	5.0	0.2	9.7
2001-2003	8	8.6	2.6	14.6	8	7.3	2.1	12.6	7	10.1	2.5	17.6	6	6.6	1.0	12.3
2002-2004	10	10.6	4.0	17.3	7	6.2	1.4	10.9	8	11.4	3.5	19.4	9	10.2	3.2	17.3
2003-2005	12	12.4	5.3	19.5	<5	3.8	0	7.7	7	10.2	2.6	17.9	6	7.5	1.4	13.7
2004-2006	13	12.7	5.7	19.8	<5	3.4	0	6.9	6	8.6	1.7	15.4	5	6.3	0.7	11.8
2005-2007	12	11.5	4.9	18.0	8	6.5	1.9	11.2	<5	5.5	0.1	10.9	<5	4.0	0	8.5
2006-2008	13	11.6	5.2	17.9	12	9.9	4.1	15.6	6	7.0	1.3	12.7	<5	5.2	0.1	10.2
2007-2009	10	8.8	3.3	14.3	11	9.1	3.6	14.6	11	13.1	5.3	20.9	8	9.4	2.8	16.0
2008-2010	7	6.1	1.5	10.6	9	7.6	2.6	12.6	12	14.3	6.1	22.5	7	7.9	2.0	13.8
2009-2011	5	4.5	0.5	8.5	<5	2.4	0	5.1	9	11.3	3.9	18.7	6	6.6	1.3	11.9

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	61.7	62.7	61.6	65.0
Median	63	64	64	67
5th percentile	31	32	43	47
95th percentile	85	86	78	76

Brain and CNS Cancer (C70-72) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	<5	2.9	0	6.3	<5	3.1	0	6.2	5	6.9	0.8	13.0	<5	5.6	0.1	11.2
2002-2004	5	5.4	0.6	10.2	<5	3.0	0	6.1	7	9.9	2.5	17.3	<5	5.6	0.1	11.2
2003-2005	9	9.4	3.2	15.5	6	5.0	0.9	9.1	6	8.7	1.7	15.8	6	7.9	1.5	14.3
2004-2006	11	10.8	4.3	17.2	5	4.1	0.4	7.8	<5	4.4	0	9.5	<5	5.0	0.0	9.9
2005-2007	11	10.1	4.1	16.1	7	5.9	1.4	10.4	<5	5.5	0.1	10.9	6	7.6	1.4	13.8
2006-2008	8	6.9	2.1	11.7	<5	2.2	0	4.9	<5	5.3	0.0	10.5	5	6.9	0.8	12.9
2007-2009	7	6.1	1.5	10.6	6	5.1	1.0	9.2	6	7.6	1.4	13.7	8	10.0	3.1	17.0
2008-2010	7	5.8	1.5	10.2	8	6.1	1.8	10.4	6	7.0	1.4	12.7	9	9.8	3.3	16.4
2009-2011	8	6.8	2.0	11.6	9	6.9	2.3	11.6	5	5.8	0.7	10.9	7	7.1	1.7	12.5

1-year survival from brain and CNS cancer in males is rising everywhere except Jersey – where it is unchanged. 5-year survival is unchanged in Guernsey, falling in Jersey, and rising in the South West and England.


1-year survival from brain and CNS cancer in females is falling in Guernsey and Jersey, and rising in the South West and England. 5-year survival is falling in Guernsey, Jersey and the South West, and rising in England.

1-year and 5-year survival for brain and CNS cancer in males is similar to England in Guernsey, Jersey and the South West.


1-year and 5-year survival for brain and CNS cancer in females is similar to England in Guernsey, Jersey and the South West.

Breast Cancer (females) (C50) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	N/A	N/A	N/A	N/A	193	172.3	147.4	197.2	N/A	N/A	N/A	N/A	116	149.6	121.2	178.0
2000-2002	N/A	N/A	N/A	N/A	220	192.3	166.2	218.5	N/A	N/A	N/A	N/A	125	157.6	128.7	186.5
2001-2003	N/A	N/A	N/A	N/A	213	181.3	156.1	206.4	N/A	N/A	N/A	N/A	137	172.2	142.0	202.4
2002-2004	N/A	N/A	N/A	N/A	201	166.8	142.9	190.7	N/A	N/A	N/A	N/A	158	195.1	163.1	227.1
2003-2005	N/A	N/A	N/A	N/A	190	157.6	134.4	180.9	N/A	N/A	N/A	N/A	164	205.0	172.1	238.0
2004-2006	N/A	N/A	N/A	N/A	193	160.3	136.9	183.7	N/A	N/A	N/A	N/A	163	202.3	170.1	234.5
2005-2007	N/A	N/A	N/A	N/A	203	165.5	142.0	189.0	N/A	N/A	N/A	N/A	134	163.0	134.5	191.5
2006-2008	N/A	N/A	N/A	N/A	212	166.7	143.5	189.9	N/A	N/A	N/A	N/A	129	155.0	127.4	182.5
2007-2009	N/A	N/A	N/A	N/A	231	178.2	154.5	201.8	N/A	N/A	N/A	N/A	121	137.6	112.1	163.1
2008-2010	N/A	N/A	N/A	N/A	236	182.3	158.4	206.2	N/A	N/A	N/A	N/A	136	156.6	129.3	183.8
2009-2011	N/A	N/A	N/A	N/A	236	185.3	161.1	209.4	N/A	N/A	N/A	N/A	145	163.5	136.0	191.1

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	63.1	64.2	62.0	62.1
Median	63	64	61	63
5th percentile	40	41	41	39
95th percentile	87	88	85	86

Breast Cancer (females) (C50) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	N/A	N/A	N/A	N/A	41	32.0	21.8	42.2	N/A	N/A	N/A	N/A	28	29.3	17.7	40.9
2002-2004	N/A	N/A	N/A	N/A	54	41.1	29.6	52.6	N/A	N/A	N/A	N/A	25	24.7	14.2	35.2
2003-2005	N/A	N/A	N/A	N/A	53	41.5	29.9	53.1	N/A	N/A	N/A	N/A	33	34.5	21.7	47.3
2004-2006	N/A	N/A	N/A	N/A	50	37.9	26.9	48.9	N/A	N/A	N/A	N/A	34	34.8	22.3	47.3
2005-2007	N/A	N/A	N/A	N/A	49	37.4	26.5	48.3	N/A	N/A	N/A	N/A	34	35.9	23.2	48.6
2006-2008	N/A	N/A	N/A	N/A	42	30.7	21.0	40.4	N/A	N/A	N/A	N/A	26	26.8	16.0	37.7
2007-2009	N/A	N/A	N/A	N/A	43	30.8	21.2	40.4	N/A	N/A	N/A	N/A	28	27.0	16.2	37.9
2008-2010	N/A	N/A	N/A	N/A	50	33.4	23.7	43.2	N/A	N/A	N/A	N/A	25	21.8	12.5	31.1
2009-2011	N/A	N/A	N/A	N/A	55	34.2	24.6	43.8	N/A	N/A	N/A	N/A	21	18.5	9.9	27.2

1-year survival from breast cancer in females is unchanged in Guernsey, and rising in Jersey, the South West and England.


5-year survival from breast cancer in females is unchanged in Guernsey and Jersey, and rising in the South West and England.

1-year survival for breast cancer in females is similar to England in Guernsey, Jersey and the South West.


5-year survival for breast cancer in females is similar to England in Guernsey and Jersey, but higher than England in the South West.

Colorectal Cancer (C18-C21) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	60	58.5	43.6	73.4	47	33.2	23.2	43.2	65	86.8	65.4	108.1	55	55.2	39.4	71.0
2000-2002	66	64.1	48.6	79.7	48	33.8	23.6	44.0	56	77.1	56.7	97.5	61	61.0	44.5	77.6
2001-2003	72	69.5	53.4	85.6	60	45.3	33.2	57.5	54	74.8	54.7	95.0	50	52.0	36.5	67.4
2002-2004	71	66.7	51.1	82.4	75	57.5	43.8	71.2	66	91.3	69.0	113.5	44	41.0	27.9	54.2
2003-2005	81	72.9	56.9	89.0	74	52.9	40.2	65.7	65	89.9	67.9	112.0	52	52.2	36.8	67.6
2004-2006	97	86.1	68.7	103.4	77	50.9	38.8	63.0	73	96.9	74.5	119.2	49	45.0	31.3	58.8
2005-2007	110	95.5	77.5	113.6	77	51.0	38.9	63.0	78	98.4	76.4	120.3	49	49.9	35.1	64.8
2006-2008	106	90.5	73.1	107.9	76	51.8	39.5	64.0	80	95.7	74.5	116.8	40	38.7	25.9	51.4
2007-2009	95	79.8	63.6	96.1	74	51.4	39.1	63.7	83	95.5	74.8	116.3	45	44.8	30.9	58.7
2008-2010	97	80.6	64.4	96.8	75	52.8	40.4	65.3	74	83.4	64.2	102.7	41	37.8	25.4	50.3
2009-2011	97	79.4	63.4	95.3	69	47.8	36.0	59.5	72	81.3	62.3	100.3	48	44.8	31.2	58.4

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	71.2	71.9	70.0	70.2
Median	73	73	70	72
5th percentile	49	50	50	50
95th percentile	88	89	88	87

Colorectal Cancer (C18-C21) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	22	20.9	12.1	29.7	18	12.6	6.4	18.9	37	48.7	32.8	64.7	24	20.2	11.2	29.2
2002-2004	32	29.7	19.3	40.1	23	16.2	9.2	23.2	27	34.2	21.1	47.2	28	21.2	12.4	30.0
2003-2005	32	29.3	19.0	39.6	23	15.7	8.9	22.6	18	22.7	12.1	33.4	26	20.0	11.3	28.7
2004-2006	31	27.7	17.8	37.6	24	16.2	9.3	23.1	21	27.0	15.4	38.6	21	16.4	8.6	24.2
2005-2007	30	25.8	16.5	35.1	28	18.6	11.3	25.9	25	31.1	18.8	43.5	15	11.6	5.2	18.1
2006-2008	24	19.9	11.8	28.0	23	15.1	8.6	21.6	22	26.3	15.2	37.5	13	10.7	4.2	17.2
2007-2009	24	18.9	11.2	26.6	24	15.3	8.9	21.7	20	22.1	12.2	32.0	15	11.6	5.0	18.1
2008-2010	33	26.2	17.2	35.3	21	12.7	7.0	18.5	19	19.9	10.8	28.9	13	9.5	3.7	15.3
2009-2011	39	30.4	20.7	40.1	19	11.9	6.2	17.5	28	29.5	18.5	40.6	16	11.7	5.1	18.2

1-year survival from colorectal cancer in males is rising everywhere. 5-year survival is unchanged in Guernsey, and rising in Jersey, the South West and England.


1-year survival from colorectal cancer in females is unchanged in Guernsey and Jersey, and rising in the South West and England. 5-year survival is falling in Guernsey, and rising in Jersey, the South West, and England.

1-year survival for colorectal cancer in males is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey and Jersey, but higher in the South West.


1-year survival for colorectal cancer in females is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey and Jersey, but higher in the South West.

Head and Neck Cancer (C00-C14, C30-C32, C73) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	45	45.2	31.9	58.5	12	11.1	4.7	17.4	20	29.2	16.3	42.1	13	15.0	6.5	23.4
2000-2002	41	41.3	28.6	54.0	20	16.6	9.0	24.3	23	34.0	20.0	48.0	14	16.5	7.5	25.4
2001-2003	46	45.5	32.3	58.7	25	21.6	12.8	30.4	31	45.6	29.5	61.8	15	18.4	8.6	28.1
2002-2004	44	43.4	30.5	56.2	29	25.1	15.6	34.5	32	47.0	30.6	63.4	12	14.3	5.8	22.7
2003-2005	52	51.3	37.3	65.3	29	25.7	16.1	35.3	37	54.3	36.7	72.0	16	21.0	10.3	31.6
2004-2006	65	62.8	47.5	78.2	27	24.6	15.1	34.1	27	38.9	24.2	53.7	15	18.8	8.9	28.7
2005-2007	68	63.5	48.3	78.7	27	23.7	14.6	32.9	25	34.3	20.8	47.7	18	22.6	11.8	33.5
2006-2008	71	64.4	49.3	79.4	22	19.0	10.9	27.1	28	36.7	23.0	50.3	15	16.7	7.7	25.6
2007-2009	62	54.8	41.0	68.5	23	18.5	10.8	26.2	31	39.6	25.6	53.7	18	20.6	10.6	30.5
2008-2010	62	53.4	40.0	66.8	20	15.6	8.5	22.6	35	43.6	29.1	58.2	15	16.7	7.9	25.6
2009-2011	59	49.3	36.6	62.0	23	18.6	10.8	26.4	33	39.8	26.1	53.5	20	21.8	11.8	31.8

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	61.8	63.2	60.1	62.1
Median	62	64	60	61
5th percentile	35	37	39	40
95th percentile	86	87	81	86

Head and Neck Cancer (C00-C14, C30-C32, C73) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	14	14.4	6.8	21.9	6	5.2	0.9	9.5	8	11.7	3.6	19.9	<5	4.7	0	9.7
2002-2004	16	16.1	8.2	24.0	6	5.2	0.9	9.5	10	13.9	5.3	22.6	5	5.8	0.4	11.2
2003-2005	17	16.9	8.9	25.0	5	4.4	0.4	8.4	11	15.6	6.3	24.8	5	6.8	0.7	12.9
2004-2006	13	12.3	5.6	19.1	<5	1.9	0	4.6	8	10.9	3.3	18.5	<5	4.2	0	8.9
2005-2007	17	16.0	8.3	23.6	5	4.6	0.6	8.7	5	6.4	0.8	12.0	<5	3.7	0	8.0
2006-2008	21	19.3	11.0	27.7	<5	2.8	0	5.9	5	6.5	0.8	12.3	<5	5.0	0.0	9.9
2007-2009	18	16.7	8.9	24.4	<5	3.6	0.1	7.2	9	11.2	3.8	18.5	<5	4.7	0.0	9.4
2008-2010	22	19.5	11.3	27.7	<5	1.8	0	4.2	10	12.4	4.7	20.1	<5	4.2	0	8.5
2009-2011	17	14.5	7.5	21.4	<5	2.5	0	5.4	8	9.7	2.9	16.5	<5	1.5	0	3.8

1-year survival from head and neck cancer in males is unchanged in Guernsey and Jersey, and rising in the South West and England. 5-year survival is falling in Guernsey, rising in Jersey, unchanged in the South West and rising in England.


1-year survival from head and neck cancer in females is unchanged in Jersey, and rising elsewhere. 5-year survival is unchanged in Guernsey, and rising elsewhere.

1-year survival for head and neck cancer in males is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.


1-year survival for head and neck cancer in females is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.

Hepatobiliary Cancer (C22-C23) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	<5	4.1	0.1	8.1	12	7.5	2.9	12.1	6	8.1	1.5	14.6	<5	4.4	0	9.4
2000-2002	<5	2.2	0	5.2	12	9.2	3.7	14.7	9	12.3	4.1	20.5	<5	4.6	0	9.4
2001-2003	<5	2.9	0	6.2	8	7.0	2.0	11.9	11	15.2	6.1	24.2	<5	2.0	0	5.0
2002-2004	7	6.6	1.7	11.6	7	5.4	1.2	9.5	9	12.6	4.3	20.9	<5	2.0	0	5.0
2003-2005	8	7.6	2.3	12.8	6	4.4	0.7	8.2	5	6.9	0.8	12.9	<5	0	0	0
2004-2006	11	10.4	4.3	16.6	6	5.3	0.9	9.6	<5	2.8	0	6.6	<5	3.5	-0.5	7.6
2005-2007	12	11.2	4.9	17.6	7	6.3	1.6	11.0	<5	2.8	0	6.6	<5	4.3	0.0	8.6
2006-2008	18	15.8	8.4	23.1	10	7.8	2.9	12.7	7	8.1	2.0	14.3	6	5.6	0.9	10.2
2007-2009	20	16.6	9.3	24.0	10	6.7	2.4	11.0	12	14.0	6.0	22.0	<5	2.1	0	4.5
2008-2010	18	14.5	7.7	21.3	10	6.2	2.1	10.3	13	14.4	6.5	22.4	<5	2.6	0	5.8
2009-2011	13	10.4	4.7	16.1	6	3.6	0.5	6.7	8	8.9	2.7	15.1	<5	2.6	0	6.3

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	71.1	71.7	71.0	70.7
Median	73	73	72	71
5th percentile	48	50	54	53
95th percentile	89	90	86	84

Hepatobiliary Cancer (C22-C23) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	<5	3.0	0	6.5	<5	1.8	0	4.3	5	6.8	0.8	12.8	<5	1.4	0	4.3
2002-2004	5	4.7	0.6	8.8	5	3.8	0.4	7.3	5	7.1	0.9	13.3	<5	0.5	0	1.6
2003-2005	6	5.4	1.1	9.7	<5	2.0	0	4.5	<5	4.2	0	8.9	<5	0.5	0	1.6
2004-2006	8	7.3	2.2	12.4	7	5.1	1.1	9.0	<5	1.3	0	3.7	<5	3.8	0	7.7
2005-2007	8	7.3	2.2	12.3	5	3.9	0.4	7.4	0	0	0	0	<5	3.2	0	7.0
2006-2008	10	9.1	3.4	14.8	7	4.8	1.1	8.6	<5	2.2	0	5.3	<5	3.2	0	6.9
2007-2009	13	10.9	4.9	16.9	7	4.8	1.1	8.6	<5	4.2	0.0	8.4	0	0	0	0
2008-2010	19	16.4	9.0	23.8	9	5.9	1.9	10.0	7	8.0	2.0	14.0	0	0	0	0
2009-2011	20	16.8	9.4	24.2	9	5.7	1.7	9.6	7	7.9	2.0	13.8	<5	2.4	0	5.6

1-year survival from hepatobiliary cancer in males is unchanged in Guernsey and Jersey, and rising in the South West and England. 5-year survival is unchanged in Jersey, and rising elsewhere.


1-year survival from hepatobiliary cancer in females is unchanged in Guernsey and Jersey, and rising in the South West and England. 5-year survival is falling in Guernsey, Jersey and the South West, and rising in England.

1-year survival for hepatobiliary cancer in males is similar to England in Guernsey, Jersey and the South West. 5-year survival is higher than England in Guernsey, and similar in Jersey and the South West.


1-year survival for hepatobiliary cancer in females is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.

Kidney and Ureter Cancer (C64-C68) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	13	12.6	5.7	19.5	5	4.4	0.5	8.3	9	12.9	4.5	21.4	7	7.6	1.6	13.6
2000-2002	11	10.6	4.3	17.0	7	6.3	1.6	11.0	9	12.2	4.2	20.3	<5	3.8	0	8.1
2001-2003	9	8.3	2.9	13.8	9	7.5	2.5	12.5	12	17.2	7.4	27.0	<5	5.7	0	11.4
2002-2004	13	12.5	5.7	19.3	17	14.1	7.2	21.0	13	19.3	8.7	29.9	12	13.3	5.4	21.3
2003-2005	18	17.2	9.2	25.2	15	11.4	5.4	17.4	18	26.4	14.1	38.7	16	18.2	8.9	27.6
2004-2006	24	22.3	13.3	31.4	14	11.1	5.0	17.1	13	17.9	8.1	27.7	18	18.2	9.4	27.1
2005-2007	20	17.2	9.6	24.9	6	4.6	0.7	8.5	12	15.8	6.8	24.8	12	11.8	4.8	18.9
2006-2008	19	16.0	8.7	23.3	5	4.6	0.6	8.7	10	12.0	4.5	19.5	9	7.4	2.2	12.7
2007-2009	23	19.2	11.2	27.1	8	5.7	1.6	9.8	21	23.2	13.1	33.2	7	5.4	1.0	9.9
2008-2010	24	20.3	12.1	28.6	10	6.7	2.4	11.0	22	22.8	13.1	32.5	6	4.9	0.6	9.2
2009-2011	21	17.7	10.1	25.4	13	8.6	3.7	13.4	19	19.3	10.5	28.1	<5	4.3	0.2	8.4

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	68.2	69.3	67.8	71.4
Median	69	70	68	74
5th percentile	45	46	48	48
95th percentile	87	88	87	87

Kidney and Ureter Cancer (C64-C68) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	<5	3.0	0	6.4	<5	0.9	0	2.6	<5	4.0	0	8.5	<5	3.0	0	6.6
2002-2004	<5	3.8	0.1	7.5	<5	0.9	0	2.6	6	8.0	1.5	14.5	<5	4.6	0	9.3
2003-2005	6	5.9	1.2	10.6	<5	2.1	0	4.6	5	6.8	0.8	12.9	<5	5.1	0.0	10.1
2004-2006	7	6.7	1.7	11.7	<5	2.7	0	5.6	6	7.8	1.5	14.1	<5	3.5	0	7.5
2005-2007	11	10.1	4.1	16.2	6	4.3	0.7	8.0	<5	3.7	0	7.9	<5	2.1	0	4.7
2006-2008	9	7.4	2.5	12.4	<5	3.1	0	6.2	<5	4.8	0.1	9.6	<5	1.0	0	2.3
2007-2009	15	12.0	5.8	18.2	<5	2.1	0	4.6	9	9.6	3.2	15.9	<5	1.9	0.0	3.7
2008-2010	13	10.5	4.7	16.3	6	2.8	0.4	5.1	11	11.4	4.5	18.2	<5	1.4	0	2.9
2009-2011	15	12.7	6.2	19.1	9	4.3	1.3	7.4	15	15.1	7.3	22.9	5	3.5	0.1	6.9

1-year survival from kidney and ureter cancer in males is unchanged in Guernsey, Jersey and the South West, and is rising in England. 5-year survival is falling in Guernsey and Jersey, unchanged in the South West, and rising in England.


1-year survival from kidney and ureter cancer in females is unchanged in Guernsey, falling in Jersey, and rising in the South West and England. 5-year survival is unchanged in Guernsey, Jersey and the South West, and rising in England.

1-year survival for kidney and ureter cancer in males is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West


1-year survival for kidney and ureter cancer in females is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.

Leukaemia (C91-C95) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	11	10.3	4.2	16.4	<5	2.9	0	5.8	13	18.2	8.2	28.2	6	6.3	0.9	11.7
2000-2002	9	8.3	2.8	13.7	<5	3.0	0	6.1	12	17.0	7.3	26.7	7	7.1	1.5	12.8
2001-2003	9	8.0	2.7	13.2	8	5.0	1.2	8.8	10	14.0	5.2	22.7	8	7.2	1.9	12.4
2002-2004	9	8.3	2.8	13.7	12	7.7	3.1	12.4	11	15.0	6.0	24.0	6	4.7	0.8	8.6
2003-2005	16	14.3	7.2	21.4	12	6.6	2.6	10.6	13	16.9	7.5	26.2	<5	3.3	0	6.7
2004-2006	16	14.2	7.1	21.3	12	7.6	3.1	12.1	9	10.6	3.6	17.7	6	4.3	0.6	8.0
2005-2007	22	18.7	10.8	26.6	12	8.0	3.2	12.9	8	9.4	2.8	16.0	8	6.1	1.6	10.6
2006-2008	30	24.8	15.8	33.8	18	13.1	6.7	19.4	12	14.5	6.2	22.7	10	7.5	2.5	12.5
2007-2009	37	30.1	20.3	40.0	18	12.0	6.1	17.9	18	21.1	11.3	31.0	10	7.9	2.7	13.2
2008-2010	38	30.6	20.7	40.4	14	9.2	4.1	14.3	20	22.4	12.4	32.3	8	6.0	1.5	10.6
2009-2011	25	20.7	12.5	28.8	8	5.2	1.4	9.0	20	21.4	11.9	30.9	14	12.0	5.3	18.7

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	69	70.7	69.5	73.9
Median	72	73	69	74
5th percentile	38	42	46	53
95th percentile	89	90	89	91

Leukaemia (C91-C95) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	8	7.8	2.4	13.1	<5	2.7	0	5.7	11	15.3	6.2	24.4	<5	2.5	0	5.6
2002-2004	8	7.6	2.3	12.9	5	3.7	0.2	7.2	7	10.0	2.5	17.5	<5	1.1	0	2.6
2003-2005	6	5.3	1.0	9.7	<5	2.6	0	5.4	11	15.0	6.0	24.1	<5	0.5	0	1.6
2004-2006	9	7.5	2.5	12.4	7	4.8	1.0	8.5	10	12.1	4.5	19.8	<5	0.5	0	1.6
2005-2007	9	7.4	2.5	12.2	7	4.2	0.9	7.5	9	10.7	3.6	17.8	<5	1.3	0	3.2
2006-2008	14	11.5	5.4	17.6	7	4.2	0.9	7.6	5	5.7	0.7	10.8	<5	1.3	0	3.1
2007-2009	12	10.1	4.3	15.8	6	3.0	0.4	5.6	7	8.2	2.1	14.2	5	3.6	0.3	6.9
2008-2010	13	10.2	4.6	15.7	7	3.4	0.7	6.1	8	9.1	2.7	15.4	5	3.6	0.3	6.8
2009-2011	7	5.5	1.4	9.6	9	5.1	1.5	8.7	7	7.8	2.0	13.6	10	6.7	2.3	11.2

1-year survival from leukaemia in males is unchanged in all areas. 5-year survival is falling in Guernsey, and rising elsewhere.


1-year survival from leukaemia in females is unchanged in Guernsey, Jersey and the South West, and rising in England. 5-year survival is unchanged in all areas.

1-year survival for leukaemia in males is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West


1-year survival for leukaemia in females is similar to England in Guernsey and the South West, but lower in Jersey. 5-year survival is similar to England in Guernsey, Jersey and the South West.

Lung Cancer (C33-C34) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	88	85.9	67.8	103.9	76	61.2	46.9	75.5	52	69.6	50.4	88.7	33	35.8	22.8	48.8
2000-2002	77	75.4	58.5	92.4	54	44.3	32.1	56.5	56	75.4	55.4	95.5	30	31.0	19.2	42.9
2001-2003	87	82.8	65.3	100.3	50	42.0	30.0	53.9	58	81.3	60.2	102.4	28	30.0	18.0	41.9
2002-2004	95	89.2	71.1	107.3	60	51.3	38.0	64.6	77	106.1	82.2	130.1	41	40.1	26.9	53.2
2003-2005	108	98.8	80.0	117.7	71	58.3	44.3	72.3	77	105.5	81.7	129.3	56	57.2	41.2	73.1
2004-2006	115	103.1	84.0	122.1	75	58.0	44.4	71.6	74	96.0	73.9	118.0	55	52.8	37.8	67.7
2005-2007	122	104.8	86.0	123.7	66	47.9	35.9	60.0	64	80.2	60.3	100.0	44	44.1	30.3	58.0
2006-2008	118	99.8	81.6	118.0	64	45.4	33.8	56.9	73	85.7	65.8	105.5	41	39.3	26.5	52.0
2007-2009	97	79.0	63.1	95.0	60	43.3	32.0	54.6	72	81.4	62.4	100.5	50	50.1	35.5	64.7
2008-2010	91	74.8	59.2	90.3	69	50.9	38.6	63.3	67	73.5	55.6	91.4	51	52.3	37.3	67.3
2009-2011	84	68.7	53.8	83.5	75	56.0	43.0	69.0	57	62.7	46.1	79.2	52	54.5	39.2	69.8

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	71.8	72.4	69.5	70.8
Median	73	73	70	71
5th percentile	53	53	51	57
95th percentile	88	89	85	85

Lung Cancer (C33-C34) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	51	48.9	35.4	62.4	26	21.4	12.9	29.9	53	70.7	51.4	89.9	27	24.5	14.5	34.4
2002-2004	65	61.7	46.6	76.8	39	30.8	20.8	40.8	55	75.3	55.2	95.4	32	28.4	17.8	39.0
2003-2005	73	67.3	51.7	82.9	41	32.1	21.8	42.4	67	90.4	68.5	112.3	37	34.0	22.2	45.8
2004-2006	78	70.2	54.4	85.9	57	43.5	31.7	55.3	66	84.8	64.1	105.4	44	39.3	26.8	51.9
2005-2007	95	81.9	65.3	98.6	59	45.8	33.7	57.9	60	73.5	54.8	92.3	40	34.4	22.9	45.9
2006-2008	88	73.5	58.0	89.1	52	39.7	28.5	50.8	59	67.7	50.2	85.2	41	35.1	23.6	46.7
2007-2009	85	69.1	54.2	84.0	42	32.7	22.6	42.8	64	71.2	53.5	88.9	35	31.8	20.5	43.0
2008-2010	78	63.3	49.0	77.5	52	36.7	26.3	47.0	67	70.7	53.5	87.9	38	37.2	24.8	49.7
2009-2011	93	74.8	59.4	90.2	69	48.3	36.5	60.2	62	64.8	48.5	81.2	40	38.7	26.1	51.4

1-year survival from lung cancer in males is unchanged in Jersey and rising in all other areas. 5-year survival is unchanged in Guernsey and the South West, and rising in Jersey and in England.


1-year survival from lung cancer in females is rising in Guernsey, the South West, and England, but falling in Jersey. 5-year survival is unchanged in Guernsey and Jersey and rising in the South West and England.

1-year survival for lung cancer in males is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, and the South West, and higher than England in Jersey.


1-year survival for lung cancer in females is higher than England in Guernsey, and similar to England in Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.

Lymphoma (C81-C85, C96) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	27	26.8	16.7	37.0	17	14.5	7.4	21.5	17	24.1	12.5	35.6	19	23.6	12.6	34.6
2000-2002	20	19.5	10.9	28.0	12	9.8	4.1	15.5	23	32.1	18.9	45.4	20	25.9	14.2	37.7
2001-2003	15	14.1	6.9	21.3	13	10.9	4.8	17.0	21	29.9	17.0	42.8	24	27.3	15.8	38.8
2002-2004	18	16.5	8.8	24.2	19	16.3	8.8	23.8	21	29.6	16.8	42.4	25	25.9	15.0	36.8
2003-2005	21	19.7	11.2	28.2	30	25.9	16.3	35.5	20	28.1	15.6	40.6	23	22.7	12.7	32.7
2004-2006	26	24.6	15.1	34.2	31	26.3	16.7	36.0	23	30.3	17.8	42.9	20	19.9	10.5	29.3
2005-2007	31	28.6	18.5	38.8	32	25.5	16.3	34.7	29	37.0	23.3	50.6	15	15.5	7.0	23.9
2006-2008	37	33.1	22.3	43.8	27	20.8	12.7	28.9	22	26.1	15.1	37.2	17	19.3	9.7	29.0
2007-2009	36	30.9	20.7	41.1	25	18.5	11.0	25.9	28	32.3	20.2	44.4	26	30.6	18.4	42.8
2008-2010	32	26.8	17.5	36.2	19	14.2	7.6	20.8	23	24.9	14.6	35.2	29	32.7	20.2	45.1
2009-2011	34	28.1	18.5	37.6	19	15.0	8.1	21.9	26	28.6	17.4	39.9	30	31.6	19.7	43.5

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	64.5	66.5	62.3	65.1
Median	67	69	64	68
5th percentile	31	34	34	37
95th percentile	87	88	85	88

Lymphoma (C81-C85, C96) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	<5	3.1	0	6.6	5	4.3	0.4	8.2	8	9.9	3.0	16.9	13	15.1	6.2	23.9
2002-2004	7	6.8	1.8	11.9	7	6.2	1.5	10.9	6	7.7	1.5	13.9	12	12.3	4.8	19.7
2003-2005	7	6.4	1.6	11.2	8	6.5	1.8	11.3	6	7.5	1.5	13.6	10	8.1	2.7	13.5
2004-2006	10	8.7	3.3	14.1	7	5.7	1.3	10.0	5	6.3	0.8	11.9	10	6.8	2.3	11.2
2005-2007	7	5.6	1.4	9.8	9	6.7	2.2	11.3	<5	4.8	0.1	9.5	8	4.7	1.4	8.0
2006-2008	10	8.3	3.1	13.6	7	5.3	1.3	9.4	6	7.1	1.4	12.9	7	4.0	1.0	7.1
2007-2009	13	10.6	4.8	16.5	9	6.5	2.0	11.0	5	6.2	0.7	11.8	<5	2.5	0.0	5.1
2008-2010	13	10.5	4.7	16.3	8	5.4	1.4	9.4	7	8.3	2.1	14.5	<5	2.2	0.0	4.4
2009-2011	14	11.1	5.2	17.0	8	5.2	1.4	9.0	5	5.7	0.7	10.7	<5	2.6	0	6.2

1-year survival from lymphoma in males is unchanged in Jersey and rising in all other areas. 5-year survival is unchanged in Jersey and rising in all other areas.


1-year survival from lymphoma in females is unchanged in Guernsey and rising in all other areas. 5-year survival is rising in all areas.

1-year survival for lymphoma in males is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.


1-year survival for lymphoma in females is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.

Male Urogenital Cancer excl. Prostate (C60-C63 excl. C61) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	15	14.9	7.2	22.5	N/A	N/A	N/A	N/A	7	10.1	2.6	17.5	N/A	N/A	N/A	N/A
2000-2002	19	19.1	10.4	27.9	N/A	N/A	N/A	N/A	8	11.8	3.6	20.1	N/A	N/A	N/A	N/A
2001-2003	22	21.6	12.5	30.8	N/A	N/A	N/A	N/A	10	14.6	5.5	23.8	N/A	N/A	N/A	N/A
2002-2004	23	23.1	13.5	32.6	N/A	N/A	N/A	N/A	10	14.7	5.5	23.9	N/A	N/A	N/A	N/A
2003-2005	23	22.5	13.2	31.8	N/A	N/A	N/A	N/A	9	13.4	4.5	22.3	N/A	N/A	N/A	N/A
2004-2006	22	21.0	12.1	29.8	N/A	N/A	N/A	N/A	7	10.7	2.8	18.6	N/A	N/A	N/A	N/A
2005-2007	18	16.6	8.9	24.4	N/A	N/A	N/A	N/A	10	15.0	5.7	24.3	N/A	N/A	N/A	N/A
2006-2008	16	14.6	7.4	21.7	N/A	N/A	N/A	N/A	8	11.5	3.5	19.5	N/A	N/A	N/A	N/A
2007-2009	10	9.0	3.4	14.6	N/A	N/A	N/A	N/A	10	14.4	5.4	23.4	N/A	N/A	N/A	N/A
2008-2010	9	8.3	2.8	13.7	N/A	N/A	N/A	N/A	8	11.7	3.6	19.9	N/A	N/A	N/A	N/A
2009-2011	16	14.5	7.4	21.6	N/A	N/A	N/A	N/A	8	11.3	3.4	19.3	N/A	N/A	N/A	N/A

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	43.9	45.5	41.1	42.3
Median	40	41	36	39
5th percentile	23	23	27	27
95th percentile	79	79	67	62

Male Urogenital Cancer excl. Prostate (C60-C63 excl. C61) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	0	0	0	0	N/A	N/A	N/A	N/A	0	0	0	0	N/A	N/A	N/A	N/A
2002-2004	0	0	0	0	N/A	N/A	N/A	N/A	0	0	0	0	N/A	N/A	N/A	N/A
2003-2005	0	0	0	0	N/A	N/A	N/A	N/A	0	0	0	0	N/A	N/A	N/A	N/A
2004-2006	<5	0.6	0	1.8	N/A	N/A	N/A	N/A	0	0	0	0	N/A	N/A	N/A	N/A
2005-2007	<5	0.6	0	1.8	N/A	N/A	N/A	N/A	0	0	0	0	N/A	N/A	N/A	N/A
2006-2008	<5	0.6	0	1.7	N/A	N/A	N/A	N/A	0	0	0	0	N/A	N/A	N/A	N/A
2007-2009	0	0	0	0	N/A	N/A	N/A	N/A	0	0	0	0	N/A	N/A	N/A	N/A
2008-2010	<5	1.0	0	2.8	N/A	N/A	N/A	N/A	0	0	0	0	N/A	N/A	N/A	N/A
2009-2011	<5	1.7	0	4.1	N/A	N/A	N/A	N/A	0	0	0	0	N/A	N/A	N/A	N/A


1-year survival from male urogenital cancers is falling in Jersey and unchanged in the South West and England. 5-year survival is unchanged in all areas.

There were no deaths in Guernsey from male urogenital cancer and so survival cannot be calculated.


1-year survival for male urogenital cancers is lower than England in Jersey and similar to England in the South West. 5-year survival is similar to England in Jersey and the South West.

Malignant Melanoma (C43) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	41	40.2	27.9	52.6	43	37.8	26.3	49.4	22	31.4	18.2	44.6	17	21.9	11.2	32.7
2000-2002	40	38.0	26.2	49.9	48	40.4	28.6	52.2	20	28.6	15.9	41.3	23	29.3	16.7	41.9
2001-2003	50	48.1	34.7	61.6	58	49.5	36.4	62.7	19	27.3	14.9	39.8	25	32.7	19.3	46.1
2002-2004	53	52.3	38.1	66.5	59	50.4	37.1	63.8	16	22.5	11.3	33.6	29	35.8	22.1	49.4
2003-2005	49	49.3	35.4	63.2	55	48.7	35.5	62.0	20	27.5	15.3	39.7	27	34.5	21.1	48.0
2004-2006	42	41.9	29.1	54.7	50	43.5	31.1	55.9	20	27.5	15.4	39.7	25	32.1	19.2	45.0
2005-2007	43	40.3	28.2	52.3	56	47.0	34.5	59.6	20	26.1	14.6	37.7	24	30.5	18.1	43.0
2006-2008	48	42.4	30.3	54.4	58	47.0	34.6	59.4	16	19.6	10.0	29.3	26	32.0	19.5	44.5
2007-2009	59	50.9	37.8	64.0	61	48.4	35.9	60.9	32	37.9	24.6	51.1	29	34.7	21.7	47.7
2008-2010	56	47.6	35.0	60.2	58	45.3	33.3	57.3	47	55.5	39.5	71.6	36	40.4	26.6	54.3
2009-2011	62	52.5	39.3	65.7	54	42.4	30.8	54.0	55	63.9	46.7	81.1	51	54.9	39.0	70.9

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	60.5	63.0	60.9	64.1
Median	62	64	62	66
5th percentile	31	34	35	38
95th percentile	86	87	86	88

Malignant Melanoma (C43) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	5	5.0	0.6	9.5	6	4.5	0.7	8.4	5	6.3	0.7	11.9	<5	3.6	0	7.8
2002-2004	5	5.1	0.6	9.5	9	5.8	1.7	9.8	6	7.8	1.5	14.2	<5	2.3	0	5.5
2003-2005	5	5.4	0.7	10.1	7	4.9	1.1	8.8	<5	5.4	0.1	10.7	<5	1.4	0	4.2
2004-2006	6	6.0	1.2	10.7	6	3.5	0.5	6.6	<5	2.8	0	6.7	<5	2.9	0	6.5
2005-2007	9	8.2	2.8	13.7	<5	2.9	0	5.8	<5	2.6	0	6.1	<5	2.8	0	6.1
2006-2008	10	8.9	3.3	14.4	<5	2.4	0	4.9	<5	3.6	0	7.7	6	4.9	0.7	9.2
2007-2009	9	7.6	2.5	12.6	<5	2.7	0	5.5	<5	3.5	0	7.6	<5	3.5	0	7.1
2008-2010	9	7.6	2.6	12.6	5	3.3	0.3	6.3	<5	1.1	0	3.3	7	6.7	1.4	11.9
2009-2011	8	6.5	1.9	11.0	5	3.8	0.4	7.1	<5	3.1	0	6.5	<5	4.3	0	8.6

1-year survival from malignant melanoma in males is unchanged in Jersey and rising in Guernsey, the South West and England. 5-year survival is rising in all areas.


1-year survival from malignant melanoma in females is unchanged in Guernsey and rising in Jersey, the South West and England. 5-year survival is rising in Guernsey, Jersey and England, but is unchanged in the South West.

1-year survival for malignant melanoma in males is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.


1-year survival for malignant melanoma in females is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.

Other Cancers (other 'C' codes excl. C44) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	52	50.0	36.3	63.7	41	30.4	20.5	40.3	34	46.6	30.8	62.4	26	27.9	16.3	39.4
2000-2002	49	46.0	33.0	58.9	38	28.4	19.0	37.9	42	57.6	40.0	75.2	28	29.9	18.0	41.8
2001-2003	55	52.5	38.6	66.5	42	34.9	24.0	45.8	45	62.4	43.9	80.8	26	29.8	17.6	42.1
2002-2004	61	56.6	42.3	70.9	48	39.2	27.8	50.6	42	55.9	38.7	73.1	28	28.9	17.3	40.5
2003-2005	56	51.7	38.0	65.3	52	42.3	30.4	54.1	35	46.5	30.9	62.1	30	29.7	18.2	41.3
2004-2006	48	43.1	30.8	55.5	43	31.0	21.3	40.8	35	44.5	29.6	59.3	34	31.0	19.7	42.3
2005-2007	46	40.6	28.7	52.4	41	29.8	20.3	39.3	38	48.2	32.7	63.7	35	33.2	21.2	45.2
2006-2008	39	33.1	22.6	43.6	33	24.8	16.1	33.6	41	49.1	34.0	64.3	38	32.6	21.1	44.2
2007-2009	44	36.6	25.6	47.6	35	28.2	18.7	37.7	53	60.8	44.2	77.5	37	28.8	18.5	39.0
2008-2010	46	37.0	26.2	47.9	40	30.7	20.9	40.5	47	51.8	36.8	66.9	41	34.9	23.4	46.4
2009-2011	59	47.3	35.1	59.5	40	30.1	20.5	39.7	53	57.7	41.9	73.6	37	33.4	21.9	44.9

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	71	71.8	68.1	71.6
Median	73	74	69	75
5th percentile	44	46	48	48
95th percentile	90	90	86	89

Other Cancers (other 'C' codes excl. C44) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	30	29.0	18.6	39.5	25	19.9	11.8	28.1	30	40.2	25.7	54.7	26	24.2	14.1	34.3
2002-2004	34	32.8	21.7	43.9	32	25.7	16.5	34.8	34	44.1	29.1	59.1	29	25.8	15.7	35.9
2003-2005	29	27.2	17.3	37.2	30	24.1	15.2	33.0	40	52.7	36.2	69.2	33	28.8	18.2	39.4
2004-2006	26	24.1	14.8	33.5	33	25.0	16.1	33.9	34	42.1	27.8	56.4	34	30.8	19.7	41.9
2005-2007	27	24.1	14.9	33.3	34	25.5	16.6	34.5	41	49.0	33.8	64.2	38	32.6	21.2	43.9
2006-2008	29	25.0	15.8	34.2	30	21.5	13.5	29.6	31	34.0	21.8	46.1	36	29.2	18.5	39.8
2007-2009	22	18.2	10.5	26.0	28	20.1	12.3	27.8	39	42.1	28.7	55.6	45	33.2	22.3	44.1
2008-2010	28	23.4	14.7	32.2	30	19.2	11.9	26.4	40	42.1	28.8	55.4	37	27.4	17.5	37.2
2009-2011	34	27.5	18.2	36.9	40	24.7	16.6	32.9	42	42.7	29.5	55.9	37	26.5	17.0	35.9

1-year survival from other cancers in males is unchanged in Jersey and rising in Guernsey, the South West and England. 5-year survival is unchanged in Guernsey, but rising elsewhere.


1-year survival from other cancers in females is unchanged in Guernsey and rising in Jersey, the South West and England. 5-year survival is rising in Guernsey and England, but is unchanged in Jersey and the South West.

1-year survival for other cancers in males is similar to England in Guernsey and Jersey and higher in the South West. 5-year survival is similar to England in Guernsey and Jersey and higher in the South West.


1-year survival for other cancers in females is similar to England in Guernsey, but higher in Jersey and the South West. 5-year survival is similar to England in Guernsey and Jersey, but higher in the South West.

Other Gynaecological Cancers (C51-C53, C57-C58) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	N/A	N/A	N/A	N/A	21	17.2	9.5	24.8	N/A	N/A	N/A	N/A	25	32.8	19.6	46.0
2000-2002	N/A	N/A	N/A	N/A	16	12.9	6.4	19.5	N/A	N/A	N/A	N/A	21	26.3	14.7	37.9
2001-2003	N/A	N/A	N/A	N/A	17	14.4	7.3	21.4	N/A	N/A	N/A	N/A	24	30.3	17.8	42.8
2002-2004	N/A	N/A	N/A	N/A	18	15.2	7.9	22.4	N/A	N/A	N/A	N/A	22	27.0	15.3	38.7
2003-2005	N/A	N/A	N/A	N/A	25	21.5	12.8	30.2	N/A	N/A	N/A	N/A	23	29.8	17.3	42.4
2004-2006	N/A	N/A	N/A	N/A	20	16.9	9.2	24.6	N/A	N/A	N/A	N/A	18	22.1	11.5	32.7
2005-2007	N/A	N/A	N/A	N/A	23	19.7	11.4	27.9	N/A	N/A	N/A	N/A	18	21.7	11.2	32.2
2006-2008	N/A	N/A	N/A	N/A	17	13.6	6.9	20.2	N/A	N/A	N/A	N/A	18	21.2	10.9	31.5
2007-2009	N/A	N/A	N/A	N/A	21	17.2	9.6	24.7	N/A	N/A	N/A	N/A	15	17.6	8.3	26.9
2008-2010	N/A	N/A	N/A	N/A	18	15.3	8.1	22.5	N/A	N/A	N/A	N/A	16	18.4	8.9	27.9
2009-2011	N/A	N/A	N/A	N/A	20	17.2	9.5	24.8	N/A	N/A	N/A	N/A	18	21.2	10.8	31.5

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	55.9	57.3	56.0	56.9
Median	55	57	57	57
5th percentile	27	27	26	30
95th percentile	87	89	85	84

Other Gynaecological Cancers (C51-C53, C57-C58) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011


Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	N/A	N/A	N/A	N/A	<5	2.4	0	5.2	N/A	N/A	N/A	N/A	6	7.0	1.2	12.8
2002-2004	N/A	N/A	N/A	N/A	<5	2.8	0	5.7	N/A	N/A	N/A	N/A	9	9.4	3.0	15.8
2003-2005	N/A	N/A	N/A	N/A	5	3.8	0.3	7.4	N/A	N/A	N/A	N/A	7	7.2	1.6	12.8
2004-2006	N/A	N/A	N/A	N/A	7	4.7	1.0	8.5	N/A	N/A	N/A	N/A	5	4.5	0.2	8.8
2005-2007	N/A	N/A	N/A	N/A	6	4.2	0.7	7.8	N/A	N/A	N/A	N/A	<5	0.5	0	1.5
2006-2008	N/A	N/A	N/A	N/A	5	3.2	0.2	6.1	N/A	N/A	N/A	N/A	<5	0.5	0	1.4
2007-2009	N/A	N/A	N/A	N/A	<5	1.8	0	3.9	N/A	N/A	N/A	N/A	<5	1.3	0	3.1
2008-2010	N/A	N/A	N/A	N/A	<5	1.7	0	3.9	N/A	N/A	N/A	N/A	<5	1.7	0	3.7
2009-2011	N/A	N/A	N/A	N/A	5	3.0	0.2	5.9	N/A	N/A	N/A	N/A	5	4.3	0.2	8.4

1-year survival from other gynaecological cancers in females is unchanged in Jersey and the South West and rising in Guernsey and England. 5-year survival is unchanged in Jersey and the South West and rising in Guernsey and England.


1-year survival for other gynaecological cancers in females is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.

Other Skin Cancers (C44) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	372	357.7	321.1	394.3	367	286.4	255.6	317.3	161	217.1	183.2	251.0	135	130.1	106.2	154.0
2000-2002	352	337.8	302.3	373.3	353	279.5	248.9	310.2	246	333.8	291.7	375.9	212	214.7	183.2	246.3
2001-2003	370	354.9	318.5	391.3	340	271.8	241.6	302.1	361	496.2	444.5	547.9	281	292.8	255.6	330.1
2002-2004	380	360.2	323.8	396.7	351	287.3	256.0	318.6	397	545.5	491.3	599.8	330	364.2	322.0	406.5
2003-2005	420	393.9	355.9	431.9	355	284.0	253.2	314.8	372	511.0	458.5	563.5	326	364.1	321.8	406.3
2004-2006	435	399.0	361.1	436.9	379	296.0	264.9	327.1	304	401.7	356.0	447.3	297	327.9	288.4	367.5
2005-2007	449	398.9	361.6	436.2	396	291.9	261.9	322.0	313	388.5	344.9	432.1	286	296.0	259.5	332.5
2006-2008	461	396.3	359.7	432.8	426	306.5	276.1	336.9	335	395.5	352.5	438.4	293	295.3	259.4	331.3
2007-2009	469	392.5	356.5	428.4	458	325.0	293.9	356.0	425	483.8	437.1	530.5	314	305.9	269.7	342.0
2008-2010	499	413.4	376.7	450.1	457	322.1	291.3	352.8	425	474.9	429.0	520.7	340	323.8	287.0	360.6
2009-2011	514	421.9	385.0	458.8	470	328.4	297.4	359.3	417	456.9	412.4	501.5	342	322.3	285.7	358.8

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	71	71.4	68.0	70.2
Median	73	73	69	72
5th percentile	45	46	43	46
95th percentile	90	90	88	89

Other Skin Cancers (C44) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011




Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	<5	4.2	0.1	8.3	<5	0.7	0	2.0	<5	3.3	0	7.1	<5	1.1	0	2.7
2002-2004	5	5.0	0.6	9.4	<5	0.7	0	2.0	<5	2.3	0	5.6	<5	2.1	0	5.4
2003-2005	<5	3.0	0	6.4	<5	0.7	0	2.0	<5	1.0	0	3.0	<5	2.7	0	6.1
2004-2006	<5	0.8	0	2.5	<5	0.7	0	1.9	<5	0.9	0	2.7	<5	2.1	0	5.3
2005-2007	<5	3.5	0.1	6.9	<5	1.0	0	2.5	<5	2.0	0	4.9	<5	0.5	0	1.5
2006-2008	5	4.1	0.5	7.8	<5	0.4	0	1.1	<5	2.2	0	5.2	<5	1.0	0	2.3
2007-2009	5	3.9	0.5	7.3	<5	0.8	0	1.9	<5	2.1	0	5.0	<5	0.9	0	2.2
2008-2010	<5	0.8	0	2.5	<5	0.4	0	1.2	<5	1.0	0	3.0	<5	2.1	0	4.3
2009-2011	0	0	0	0	<5	1.2	0	2.8	<5	1.7	0	4.1	<5	1.7	0	3.7


Due to the very small numbers of deaths from other skin cancers, survival cannot be calculated.

Ovarian Cancer (C56) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	N/A	N/A	N/A	N/A	27	23.7	14.6	32.7	N/A	N/A	N/A	N/A	24	31.4	18.4	44.5
2000-2002	N/A	N/A	N/A	N/A	23	20.4	11.9	28.8	N/A	N/A	N/A	N/A	27	32.9	19.8	45.9
2001-2003	N/A	N/A	N/A	N/A	22	19.1	11.0	27.2	N/A	N/A	N/A	N/A	34	40.9	26.4	55.4
2002-2004	N/A	N/A	N/A	N/A	25	21.5	12.9	30.2	N/A	N/A	N/A	N/A	30	33.8	21.0	46.6
2003-2005	N/A	N/A	N/A	N/A	23	20.1	11.7	28.5	N/A	N/A	N/A	N/A	27	31.1	18.9	43.4
2004-2006	N/A	N/A	N/A	N/A	32	26.8	17.3	36.3	N/A	N/A	N/A	N/A	19	22.6	12.2	33.0
2005-2007	N/A	N/A	N/A	N/A	34	27.6	18.0	37.2	N/A	N/A	N/A	N/A	23	27.4	15.8	38.9
2006-2008	N/A	N/A	N/A	N/A	36	27.5	18.2	36.8	N/A	N/A	N/A	N/A	24	26.4	15.3	37.5
2007-2009	N/A	N/A	N/A	N/A	26	18.5	11.1	26.0	N/A	N/A	N/A	N/A	22	21.8	12.1	31.5
2008-2010	N/A	N/A	N/A	N/A	27	19.2	11.7	26.6	N/A	N/A	N/A	N/A	18	17.5	8.9	26.0
2009-2011	N/A	N/A	N/A	N/A	22	15.9	9.1	22.8	N/A	N/A	N/A	N/A	12	11.6	4.7	18.6

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	64.7	66.2	65.9	67.4
Median	66	68	71	72
5th percentile	37	38	41	40
95th percentile	87	88	90	85

Ovarian Cancer (C56) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011


Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	N/A	N/A	N/A	N/A	6	5.3	0.9	9.6	N/A	N/A	N/A	N/A	19	18.3	9.4	27.3
2002-2004	N/A	N/A	N/A	N/A	8	7.3	2.1	12.4	N/A	N/A	N/A	N/A	27	28.8	17.1	40.5
2003-2005	N/A	N/A	N/A	N/A	10	8.9	3.2	14.6	N/A	N/A	N/A	N/A	25	29.0	17.0	40.9
2004-2006	N/A	N/A	N/A	N/A	11	8.4	3.2	13.6	N/A	N/A	N/A	N/A	20	23.6	12.9	34.3
2005-2007	N/A	N/A	N/A	N/A	11	7.3	2.7	11.8	N/A	N/A	N/A	N/A	13	13.7	6.1	21.3
2006-2008	N/A	N/A	N/A	N/A	11	6.6	2.4	10.8	N/A	N/A	N/A	N/A	18	17.8	9.2	26.5
2007-2009	N/A	N/A	N/A	N/A	15	9.6	4.5	14.7	N/A	N/A	N/A	N/A	19	18.0	9.2	26.7
2008-2010	N/A	N/A	N/A	N/A	21	13.0	7.1	19.0	N/A	N/A	N/A	N/A	18	16.6	8.2	25.0
2009-2011	N/A	N/A	N/A	N/A	26	17.1	10.2	24.0	N/A	N/A	N/A	N/A	12	10.1	3.7	16.6

1-year survival from ovarian cancer in females is rising in Guernsey and England, falling in Jersey and unchanged in the South West. 5-year survival is unchanged in Guernsey, Jersey and the South West but is rising in England.


1-year survival for ovarian cancer in females is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey and the South West, but is higher in Jersey than in England.

Prostate Cancer (C61) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	102	95.7	77.0	114.4	N/A	N/A	N/A	N/A	79	107.6	83.7	131.5	N/A	N/A	N/A	N/A
2000-2002	123	115.5	95.0	136.1	N/A	N/A	N/A	N/A	89	121.2	95.8	146.6	N/A	N/A	N/A	N/A
2001-2003	161	153.5	129.6	177.3	N/A	N/A	N/A	N/A	98	133.8	107.1	160.5	N/A	N/A	N/A	N/A
2002-2004	215	203.0	175.7	230.3	N/A	N/A	N/A	N/A	97	133.0	106.3	159.8	N/A	N/A	N/A	N/A
2003-2005	218	204.4	177.0	231.7	N/A	N/A	N/A	N/A	100	138.5	111.1	165.9	N/A	N/A	N/A	N/A
2004-2006	227	209.1	181.7	236.6	N/A	N/A	N/A	N/A	90	118.4	93.7	143.0	N/A	N/A	N/A	N/A
2005-2007	206	183.7	158.4	209.0	N/A	N/A	N/A	N/A	103	127.7	102.8	152.7	N/A	N/A	N/A	N/A
2006-2008	216	182.7	158.0	207.3	N/A	N/A	N/A	N/A	111	126.3	102.5	150.2	N/A	N/A	N/A	N/A
2007-2009	188	153.2	131.1	175.4	N/A	N/A	N/A	N/A	134	150.6	124.7	176.4	N/A	N/A	N/A	N/A
2008-2010	206	168.9	145.5	192.2	N/A	N/A	N/A	N/A	133	148.0	122.5	173.6	N/A	N/A	N/A	N/A
2009-2011	219	183.5	159.0	208.1	N/A	N/A	N/A	N/A	115	128.6	104.8	152.4	N/A	N/A	N/A	N/A

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	71.3	72.1	69.5	70.2
Median	71	72	70	70
5th percentile	56	57	54	53
95th percentile	87	88	84	87

Prostate Cancer (C61) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011


Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	20	17.9	10.0	25.8	N/A	N/A	N/A	N/A	36	44.7	30.0	59.5	N/A	N/A	N/A	N/A
2002-2004	30	27.5	17.6	37.4	N/A	N/A	N/A	N/A	30	38.1	24.3	51.8	N/A	N/A	N/A	N/A
2003-2005	33	29.7	19.4	39.9	N/A	N/A	N/A	N/A	31	38.0	24.5	51.5	N/A	N/A	N/A	N/A
2004-2006	37	32.4	21.8	42.9	N/A	N/A	N/A	N/A	27	32.3	20.0	44.5	N/A	N/A	N/A	N/A
2005-2007	43	36.3	25.3	47.3	N/A	N/A	N/A	N/A	34	38.1	25.2	51.0	N/A	N/A	N/A	N/A
2006-2008	46	36.4	25.7	47.1	N/A	N/A	N/A	N/A	31	32.6	21.0	44.2	N/A	N/A	N/A	N/A
2007-2009	47	36.8	26.1	47.5	N/A	N/A	N/A	N/A	43	43.0	30.0	56.0	N/A	N/A	N/A	N/A
2008-2010	48	36.1	25.8	46.5	N/A	N/A	N/A	N/A	46	46.0	32.5	59.4	N/A	N/A	N/A	N/A
2009-2011	46	35.4	25.0	45.8	N/A	N/A	N/A	N/A	52	51.3	37.1	65.4	N/A	N/A	N/A	N/A

1-year survival from prostate cancer in males is unchanged in Guernsey but rising in Jersey, the South West and England. 5-year survival is unchanged in Jersey but rising in Guernsey, the South West and England.


1-year survival for prostate cancer in males is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.

Upper Gastrointestinal Cancers (C15-C16, C25) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	42	37.9	26.3	49.5	39	24.5	16.3	32.7	45	62.5	44.1	81.0	27	24.7	14.5	34.9
2000-2002	41	38.7	26.8	50.7	36	22.5	14.6	30.3	46	63.0	44.6	81.4	28	26.4	15.7	37.0
2001-2003	36	34.9	23.4	46.3	32	20.4	12.8	27.9	41	55.4	38.3	72.6	25	26.2	15.1	37.3
2002-2004	44	42.8	30.1	55.5	34	23.1	14.9	31.2	47	63.4	45.0	81.7	32	31.8	19.8	43.8
2003-2005	45	42.0	29.6	54.4	41	28.2	19.1	37.3	47	63.4	45.1	81.8	33	32.4	20.5	44.3
2004-2006	64	58.1	43.7	72.5	48	34.3	24.1	44.5	45	59.0	41.6	76.5	27	25.1	14.9	35.3
2005-2007	62	54.5	40.8	68.2	49	34.1	24.1	44.1	43	54.9	38.3	71.5	32	28.8	18.0	39.5
2006-2008	67	58.3	44.2	72.3	48	31.3	22.0	40.6	45	55.1	38.8	71.4	32	26.9	16.7	37.0
2007-2009	54	45.2	33.0	57.4	35	22.2	14.4	29.9	47	57.1	40.6	73.7	39	31.1	20.4	41.8
2008-2010	51	42.3	30.6	54.0	36	23.0	15.1	30.9	45	53.3	37.5	69.0	30	23.5	14.4	32.6
2009-2011	53	43.5	31.7	55.3	36	23.1	15.1	31.0	43	49.8	34.7	64.8	28	22.7	13.4	32.0

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	72.1	73.1	70.1	70.3
Median	73	74	72	71
5th percentile	50	52	48	50
95th percentile	89	90	89	90

Upper Gastrointestinal Cancers (C15-C16, C25) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	23	22.6	13.3	31.8	22	13.6	7.6	19.6	38	51.4	34.9	67.9	20	16.1	8.4	23.8
2002-2004	28	27.2	17.1	37.3	30	18.3	11.4	25.2	42	55.3	38.4	72.3	27	23.9	13.9	33.8
2003-2005	34	32.2	21.3	43.1	31	19.8	12.4	27.2	37	48.3	32.5	64.1	31	28.4	17.5	39.3
2004-2006	40	36.8	25.4	48.3	32	21.9	13.9	30.0	38	48.3	32.8	63.8	29	26.1	15.7	36.6
2005-2007	49	43.8	31.5	56.2	31	22.1	13.9	30.3	32	38.4	24.9	51.9	27	23.9	14.1	33.8
2006-2008	50	43.4	31.3	55.6	35	23.5	15.3	31.6	30	35.6	22.6	48.5	27	22.2	13.1	31.3
2007-2009	44	36.8	25.8	47.8	31	19.9	12.5	27.2	36	40.9	27.3	54.5	33	26.5	16.7	36.3
2008-2010	49	39.6	28.4	50.9	35	21.9	14.3	29.6	35	39.2	26.1	52.4	28	19.9	11.9	27.9
2009-2011	48	38.0	27.1	48.9	27	16.4	9.8	23.0	42	45.8	31.7	59.8	26	17.6	10.2	25.0

1-year survival from upper gastrointestinal cancers in males is rising in all areas. 5-year survival is rising in all areas.


1-year survival from upper gastrointestinal cancers in females is unchanged in Jersey, but rising in all other areas. 5-year survival is rising in Jersey and England but is unchanged in Guernsey and the South West.

1-year survival for upper gastrointestinal cancers in males is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.


1-year survival for upper gastrointestinal cancers in females is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey, Jersey and the South West.

Uterine Cancer (C54-C55) - INCIDENCE


Incidence rates in males, 1999-2011




Incidence rates in females, 1999-2011



Incidence rates in males, latest five years



Incidence rates in females, latest five years



Age-standardised incidence rates (per 100,000 population), 1999-2011


Years of Diagnosis	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
1999-2001	N/A	N/A	N/A	N/A	19	16.7	8.9	24.4	N/A	N/A	N/A	N/A	12	12.3	4.8	19.7
2000-2002	N/A	N/A	N/A	N/A	19	17.2	9.3	25.1	N/A	N/A	N/A	N/A	11	10.3	3.7	17.0
2001-2003	N/A	N/A	N/A	N/A	18	15.6	8.2	23.1	N/A	N/A	N/A	N/A	16	16.5	7.7	25.3
2002-2004	N/A	N/A	N/A	N/A	17	14.5	7.4	21.6	N/A	N/A	N/A	N/A	15	16.0	7.2	24.8
2003-2005	N/A	N/A	N/A	N/A	17	14.7	7.6	21.9	N/A	N/A	N/A	N/A	19	21.1	10.9	31.3
2004-2006	N/A	N/A	N/A	N/A	16	13.6	6.8	20.3	N/A	N/A	N/A	N/A	14	15.1	6.7	23.5
2005-2007	N/A	N/A	N/A	N/A	25	18.7	11.1	26.2	N/A	N/A	N/A	N/A	18	19.2	9.8	28.7
2006-2008	N/A	N/A	N/A	N/A	25	18.2	10.8	25.6	N/A	N/A	N/A	N/A	20	22.5	12.2	32.8
2007-2009	N/A	N/A	N/A	N/A	28	20.0	12.3	27.7	N/A	N/A	N/A	N/A	26	28.6	17.0	40.2
2008-2010	N/A	N/A	N/A	N/A	23	18.5	10.7	26.2	N/A	N/A	N/A	N/A	27	29.7	18.0	41.4
2009-2011	N/A	N/A	N/A	N/A	27	20.6	12.5	28.6	N/A	N/A	N/A	N/A	23	23.8	13.5	34.0

Age at diagnosis, 2007-2011 combined data (2006-2010 for England)


	England	South West	Jersey	Guernsey
Mean	66.4	67.6	68.4	65.7
Median	66	67	68	64
5th percentile	47	49	49	50
95th percentile	86	87	88	87

Uterine Cancer (C54-C55) - MORTALITY


Mortality rates in males, 2001-2011




Mortality rates in females, 2001-2011



Mortality rates in males, 2007-2011



Mortality rates in females, 2007-2011



Age-standardised mortality rates (per 100,000 population), 2001-2011

Years of Death	Jersey								Guernsey							
	Males				Females				Males				Females			
	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI	Count	ASR	LCI	UCI
2001-2003	N/A	N/A	N/A	N/A	<5	0.4	0	1.2	N/A	N/A	N/A	N/A	<5	2.2	0.0	4.4
2002-2004	N/A	N/A	N/A	N/A	<5	0.8	0	2.0	N/A	N/A	N/A	N/A	7	4.7	0.9	8.6
2003-2005	N/A	N/A	N/A	N/A	<5	1.1	0	2.6	N/A	N/A	N/A	N/A	<5	3.1	0	6.4
2004-2006	N/A	N/A	N/A	N/A	<5	2.9	0	5.9	N/A	N/A	N/A	N/A	<5	3.0	0	6.2
2005-2007	N/A	N/A	N/A	N/A	8	5.3	1.5	9.1	N/A	N/A	N/A	N/A	<5	0.5	0	1.5
2006-2008	N/A	N/A	N/A	N/A	10	6.8	2.5	11.1	N/A	N/A	N/A	N/A	<5	1.8	0	4.6
2007-2009	N/A	N/A	N/A	N/A	9	5.4	1.7	9.1	N/A	N/A	N/A	N/A	<5	3.0	0	6.7
2008-2010	N/A	N/A	N/A	N/A	6	3.4	0.5	6.2	N/A	N/A	N/A	N/A	<5	2.5	0	5.9
2009-2011	N/A	N/A	N/A	N/A	<5	1.7	0.0	3.5	N/A	N/A	N/A	N/A	8	7.3	1.9	12.7

1-year survival from uterine cancer in females is rising in Guernsey and England, and is unchanged in Jersey and the South West. 5-year survival is rising in Guernsey, the South West and England, but is falling in Jersey.

1-year survival for uterine cancer in females is similar to England in Guernsey, Jersey and the South West. 5-year survival is similar to England in Guernsey and Jersey, and higher in the South West.

Staging completeness

Percentage of cancers with valid stage, for main five cancers, in Guernsey

Site	Year				
	2007	2008	2009	2010	2011
Breast	10%	2%	9%	0%	25%
Prostate	7%	0%	4%	2%	12%
Lung	11%	13%	21%	17%	25%
Colorectal	35%	22%	62%	8%	19%
Malignant Melanoma	36%	25%	35%	69%	74%

Percentage of cancers with valid stage, for main five cancers, in Jersey

Site	Year				
	2007	2008	2009	2010	2011
Breast	4%	5%	1%	5%	20%
Prostate	3%	3%	2%	2%	2%
Lung	19%	12%	22%	23%	15%
Colorectal	61%	54%	59%	54%	47%
Malignant Melanoma	78%	82%	77%	88%	83%

Appendix – ICD-10 codes

ICD-10 Code	Site description
C00	Malignant neoplasm of lip
C01	Malignant neoplasm of base of tongue
C02	Malignant neoplasm of other and unspecified parts of tongue
C03	Malignant neoplasm of gum
C04	Malignant neoplasm of floor of mouth
C05	Malignant neoplasm of palate
C06	Malignant neoplasm of other and unspecified parts of mouth
C07	Malignant neoplasm of parotid gland
C08	Malignant neoplasm of other and unspecified major salivary glands
C09	Malignant neoplasm of tonsil
C10	Malignant neoplasm of oropharynx
C11	Malignant neoplasm of nasopharynx
C12	Malignant neoplasm of pyriform sinus
C13	Malignant neoplasm of hypopharynx
C14	Malignant neoplasm of ill-defined sites lip/oral cavity/pharynx
C15	Malignant neoplasm of oesophagus
C16	Malignant neoplasm of stomach
C17	Malignant neoplasm of small intestine
C18	Malignant neoplasm of colon
C19	Malignant neoplasm of rectosigmoid junction
C20	Malignant neoplasm of rectum
C21	Malignant neoplasm of anus and anal canal
C22	Malignant neoplasm of liver and intrahepatic bile ducts
C23	Malignant neoplasm of gallbladder
C24	Malignant neoplasm of other and unspecified parts biliary tract
C25	Malignant neoplasm of pancreas
C26	Malignant neoplasm of other and ill-defined digestive organs
C30	Malignant neoplasm of nasal cavity and middle ear
C31	Malignant neoplasm of accessory sinuses
C32	Malignant neoplasm of larynx
C33	Malignant neoplasm of trachea
C34	Malignant neoplasm of bronchus and lung
C37	Malignant neoplasm of thymus
C38	Malignant neoplasm of heart, mediastinum and pleura
C39	Malignant neoplasm of other and ill-defined sites respiratory system + intrathoracic organs
C40	Malignant neoplasm of bone and articular cartilage of limbs
C41	Malignant neoplasm of bone + articular cartilage of other + unspecified sites
C43	Malignant melanoma of skin
C44	Other malignant neoplasms of skin
C45	Mesothelioma
C46	Kaposi's sarcoma
C47	Malignant neoplasm of peripheral nerves and autonomic nervous system
C48	Malignant neoplasm of retroperitoneum and peritoneum
C49	Malignant neoplasm of other connective and soft tissue

ICD-10 Code	Site description
C50	Malignant neoplasm of breast
C51	Malignant neoplasm of vulva
C52	Malignant neoplasm of vagina
C53	Malignant neoplasm of cervix uteri
C54	Malignant neoplasm of corpus uteri
C55	Malignant neoplasm of uterus, part unspecified
C56	Malignant neoplasm of ovary
C57	Malignant neoplasm of other and unspec female genital orgs
C58	Malignant neoplasm of placenta
C60	Malignant neoplasm of penis
C61	Malignant neoplasm of prostate
C62	Malignant neoplasm of testis
C63	Malignant neoplasm of other and unspec male genital organs
C64	Malignant neoplasm of kidney, except renal pelvis
C65	Malignant neoplasm of renal pelvis
C66	Malignant neoplasm of ureter
C67	Malignant neoplasm of bladder
C68	Malignant neoplasm of other and unspecified urinary organs
C69	Malignant neoplasm of eye and adnexa
C70	Malignant neoplasm of meninges
C71	Malignant neoplasm of brain
C72	Malig neopl spinal cord, cranial nerves & oth parts of CNS
C73	Malignant neoplasm of thyroid gland
C74	Malignant neoplasm of adrenal gland
C75	Malignant neo other endocrine glands and related structures
C76	Malignant neoplasm of other and ill-defined sites
C77	Secondary and unspecified malignant neoplasm of lymph nodes
C78	Sec malignant neoplasm of respiratory and digestive organs
C79	Secondary malignant neoplasm of other sites
C80	Malignant neoplasm without specification of site
C81	Hodgkin's disease
C82	Follicular [nodular] non-Hodgkin's lymphoma
C83	Diffuse non-Hodgkin's lymphoma
C84	Peripheral and cutaneous T-cell lymphomas
C85	Other and unspecified types of non-Hodgkin's lymphoma
C88	Malignant immunoproliferative diseases
C90	Multiple myeloma and malignant plasma cell neoplasms
C91	Lymphoid leukaemia
C92	Myeloid leukaemia
C93	Monocytic leukaemia
C94	Other leukaemias of specified cell type
C95	Leukaemia of unspecified cell type
C96	Oth & unspec malig neop lymphoid, haematopoietic & rel tiss
C97	Malignant neoplasms of independent (primary) multiple sites